

GROWTH OF BALLS OF HOLOMORPHIC SECTIONS AND ENERGY AT EQUILIBRIUM

ROBERT BERMAN, SÉBASTIEN BOUCKSOM

ABSTRACT. Let L be a big line bundle on a compact complex manifold X . Given a non-pluripolar compact subset K of X and the weight ϕ of a continuous Hermitian metric $e^{-\phi}$ on L , we define the energy at equilibrium of (K, ϕ) as the Aubin-Mabuchi energy of the extremal psh weight associated to (K, ϕ) . We prove the differentiability of the energy at equilibrium with respect to ϕ , and we show that this energy describes the asymptotic behaviour as $k \rightarrow \infty$ of the volume of the sup-norm unit ball induced by $(K, k\phi)$ on the space of global holomorphic sections $H^0(X, kL)$. As a consequence of these results, we recover and extend Rumely's Robin-type formula for the transfinite diameter. We also obtain an asymptotic description of the analytic torsion, and extend Yuan's equidistribution theorem for algebraic points of small height to the case of a big line bundle.

CONTENTS

Introduction	2
1. Mixed Monge-Ampère operators and equilibrium weights	9
2. The Bergman distortion function and the Bernstein-Markov property	15
3. The Aubin-Mabuchi energy functional	19
4. Volume growth and transfinite diameter	27
5. Applications to logarithmic pluri-potential theory	30
6. Analytic torsion and equidistribution of small points	35
References	39
Bibliography	39

INTRODUCTION

0.1. The setting. Let L be a holomorphic line bundle over a compact complex manifold X of dimension n . By a *weighted subset* (K, ϕ) (resp. a *weighted measure* (μ, ϕ)), we will mean the data of a non-pluripolar compact subset K of X (resp. a probability measure with non-pluripolar support) together with the weight ϕ of a continuous Hermitian metric $e^{-\phi}$ on the restriction $L|_K$.

Using additive notation for tensor powers, we can then endow the space of global sections $s \in H^0(X, kL)$ of kL with the L^2 -norm

$$\|s\|_{L^2(\mu, k\phi)}^2 := \int_X |s|_{k\phi}^2 d\mu$$

and the L^∞ -norm

$$\|s\|_{L^\infty(K, k\phi)} := \sup_K |s|_{k\phi}.$$

Consider the special case where K and $\text{supp } \mu$ are compact subsets of

$$\mathbb{C}^n \subset \mathbb{P}^n =: X$$

endowed with the ample line bundle $\mathcal{O}(1) =: L$. Restricting to $\mathbb{C}^n$ identifies $H^0(\mathbb{P}^n, \mathcal{O}(k))$ with the space of polynomials of total degree at most k . The section of L cutting out the hyperplane at infinity induces a (flat) Hermitian metric on L over $\mathbb{C}^n$, so that a continuous weight ϕ on $L|_K$ is naturally identified with a *function* in $C^0(K)$. On the other hand a pluri-subharmonic (psh for short) function on $\mathbb{C}^n$ with at most logarithmic growth at infinity gets identified with the weight ϕ of a non-negatively curved (singular) Hermitian metric on L , which will thus be referred to as a *psh weight*.

In the general setting described above, the asymptotic study as $k \rightarrow \infty$ of $H^0(X, kL)$ endowed with the above L^2 or L^∞ -norms thus appears as a natural generalisation of the classical theory of orthogonal polynomials (cf. for instance [STBook] and in particular Bloom's appendix therein).

These two norms on $H^0(kL)$ are equivalently described by their unit balls, which will respectively be denoted by

$$\mathcal{B}^2(\mu, k\phi), \mathcal{B}^\infty(K, k\phi) \subset H^0(kL).$$

The main goal of the present paper is to study the asymptotic behaviour of the volume of these balls as $k \rightarrow \infty$. As we shall see, it is related to a well-known energy functional that we now describe.

0.2. The Aubin-Mabuchi energy functional. We denote the curvature $(1, 1)$ -form of a smooth weight ϕ on L as $dd^c \phi$, and define the *Monge-Ampère operator* on such weights as

$$\text{MA}(\phi) := (dd^c \phi)^n.$$

We have normalised as usual the operator d^c so that $dd^c = \frac{i}{\pi} \partial \bar{\partial}$.

Integrating against this measure-valued operator induces a 1-form on the (affine) space of smooth weights on L , and it is a remarkable fact that this 1-form is *closed*, hence exact. The primitive of this Monge-Ampère 1-form will be denoted

by $\phi \mapsto \mathcal{E}(\phi)$ and called the *Aubin-Mabuchi* energy functional. It is therefore characterised by the property

$$\frac{d}{dt} \bigg|_{t=0} \mathcal{E}((1-t)\phi_1 + t\phi_2) = \int_X (\phi_2 - \phi_1) \text{MA}(\phi_1). \quad (0.1)$$

As any primitive, $\mathcal{E}$ is only defined up to a constant. We will always assume that it is normalised by $\mathcal{E}(\phi_0) = 0$ for some auxiliary weight ϕ_0 fixed once and for all. On the other hand, *differences* $\mathcal{E}(\phi) - \mathcal{E}(\psi)$ are intrinsically defined. An explicit formula for $\mathcal{E}$ can be obtained by integration along line segments, which yields

$$\mathcal{E}(\phi) - \mathcal{E}(\psi) = \frac{1}{n+1} \sum_{j=0}^n \int_X (\phi - \psi) (dd^c \phi)^j \wedge (dd^c \psi)^{n-j}. \quad (0.2)$$

Note that the right-hand side involves the Bott-Chern secondary class attached to the Chern character. The functional $\mathcal{E}$ seems to have been first explicitly mentioned in Mabuchi's paper [Mab86], where it is denoted by L . It however only differs from Aubin's J -functional [Aub84] by a trivial term, hence the chosen terminology. It also corresponds to the functional I in [Che00, Don05a] and to minus $F_{\omega_0}^0$ on p.59 of Tian's book [TiaBook], where it is proved that $\phi \mapsto \mathcal{E}(\phi)$ is non-decreasing and concave on smooth psh weights.

By the fundamental work of Bedford-Taylor, mixed Monge-Ampère operators can be extended to locally bounded psh weights ϕ . Since the difference of two such weights is a bounded function on X , we can use formula (0.2) to *define* the Aubin-Mabuchi energy $\mathcal{E}(\phi)$ for a locally bounded weight ϕ . The proofs of all the above properties, which only rely on integration by parts, are then easily extended to this setting.

The locally bounded case is good enough for our purposes when L is *ample*. The more general situation of a *big* line bundle is treated in Section 3 relying on non-pluripolar products of currents and the appropriate integration-by-part formula proved in [BEGZ08]. The end result is that $\mathcal{E}(\phi)$ defined by (0.2) for two psh weights ϕ, ψ with minimal singularities still satisfies (0.1) above. It is non-decreasing and concave, and is continuous along monotonic sequences of such weights.

0.3. Asymptotics of ball volumes and energy at equilibrium. Assume now that L is a big line bundle (which implies that X is bimeromorphic to projective manifold). Given a weighted subset (K, ϕ) , its *equilibrium weight* is defined as the following extremal weight:

$$P_K \phi := \sup^* \{ \psi \text{ psh weight, } \psi \leq \phi \text{ on } K \}, \quad (0.3)$$

where the star denotes upper semi-continuous regularisation. The equilibrium weight is itself a psh weight with minimal singularities (recall that K is assumed to be non-pluripolar throughout). The *equilibrium measure* of (K, ϕ) is the probability measure defined by

$$\mu_{\text{eq}}(K, \phi) := \text{vol}(L)^{-1} \text{MA}(P_K \phi). \quad (0.4)$$

The normalising factor is the *volume* of L , i.e.

$$\text{vol}(L) = \lim_{k \rightarrow \infty} \frac{n!}{k^n} N_k$$

where $N_k := h^0(kL)$ denotes the complex dimension of $H^0(kL)$ (cf. Theorem 1.2). Note that $\text{vol}(L) > 0$, precisely because L is big.

The measure $\mu_{\text{eq}}(K, \phi)$ is concentrated on K , and $P_K \phi = \phi$ holds a.e. on K with respect to this measure (cf. Proposition 1.10). We define the *energy at equilibrium* of (K, ϕ) as

$$\mathcal{E}_{\text{eq}}(K, \phi) := \text{vol}(L)^{-1} \mathcal{E}(P_K \phi). \quad (0.5)$$

The energy at equilibrium is well-defined only up to an overall additive constant, but *differences*

$$\mathcal{E}_{\text{eq}}(K_1, \phi_1) - \mathcal{E}_{\text{eq}}(K_2, \phi_2)$$

are intrinsically defined. Our choice of normalisation yields the scaling property

$$\mathcal{E}_{\text{eq}}(K, \phi + c) = \mathcal{E}_{\text{eq}}(K, \phi) + c \quad (0.6)$$

for each constant $c \in \mathbb{R}$.

On the other hand, for each k we introduce the $\mathcal{L}$ -functional

$$\mathcal{L}_k(K, \phi) := \frac{1}{2kN_k} \log \text{vol}_k \mathcal{B}^\infty(K, k\phi), \quad (0.7)$$

which is meant to be reminiscent of Donaldson's $\mathcal{L}$ -functionals [Don05a]. The volume vol_k denotes Lebesgue measure on the vector space $H^0(kL)$, and is thus only defined up to a multiplicative constant. As a consequence, the functional $\mathcal{L}_k$ is defined up to an overall additive constant depending on k , but here again *differences* $\mathcal{L}_k(K_1, \phi_1) - \mathcal{L}_k(K_2, \phi_2)$ are well-defined since they do not depend on the choice of vol_k . Since $H^0(kL)$ has real dimension $2N_k$, our choice of normalisation yields

$$\mathcal{L}_k(K, \phi + c) = \mathcal{L}_k(K, \phi) + c \quad (0.8)$$

for each constant $c \in \mathbb{R}$, which should of course be compared to (0.6). Equivalently $\mathcal{L}_k$ defines a single valued function of (K, ϕ) relatively to a fixed reference weighted set, if vol_k is taken as the Lebesgue measure which gives a unit mass to the corresponding reference ball.

We now describe our first main result:

Theorem A. Let X be a compact complex manifold and L be a big line bundle. For any two weighted subsets (K_1, ϕ_1) and (K_2, ϕ_2) of X we have

$$\lim_{k \rightarrow \infty} (\mathcal{L}_k(K_1, \phi_1) - \mathcal{L}_k(K_2, \phi_2)) = \mathcal{E}_{\text{eq}}(K_1, \phi_1) - \mathcal{E}_{\text{eq}}(K_2, \phi_2).$$

Our second main result is a complex analogue of a result of Alexandrov in the setting of convex geometry [Ale38], see also [SchBook] p.345.

Theorem B. Let L be a big line bundle on a compact complex manifold X , and let K be a non-pluripolar compact subset of X . Then $\phi \mapsto \mathcal{E}_{\text{eq}}(K, \phi)$ is concave and continuous on the space of continuous weights on $L|_K$ endowed with the

topology of C^0 convergence. It admits linear directional derivatives, which are given by integration against the equilibrium measure:

$$\frac{d}{dt}\bigg|_{t=0} \mathcal{E}_{\text{eq}}(K, \phi + tv) = \langle v, \mu_{\text{eq}}(K, \phi) \rangle$$

for every $v \in C^0(K)$.

The crucial assertion here is the differentiability property, which means that the super-differentials of the concave function $\phi \mapsto \mathcal{E}_{\text{eq}}(K, \phi)$ are all reduced to a point. This can be understood as a *linear response* property for the energy at equilibrium. Note however that $t \mapsto \mathcal{E}_{\text{eq}}(K, \phi + tv)$ is not *twice* differentiable in general, even for smooth ϕ and v with $K = X = \mathbb{P}^1$ (cf. Example 3.12).

Theorem B is actually a crucial ingredient of the proof of Theorem A. It is also a key tool in the proof of the arithmetic equidistribution result to be described below (Theorem D), as well as in several other equidistribution theorems [BerBou08b, BWN08, BBWN08].

This differentiability result bears a strong resemblance with the differentiability property of the volume of cohomology classes [BFJ07], which is in some sense a non-archimedean analogue of the present result (compare [BFJ08]).

0.4. Volume of L^2 -balls and transfinite diameter. Similarly to (0.7) we introduce $\mathcal{L}$ -functionals on weighted measures by

$$\mathcal{L}_k(\mu, \phi) := \frac{1}{2kN_k} \log \text{vol}_k \mathcal{B}^2(\mu, k\phi).$$

These functionals (which are actually closer to Donaldson's $\mathcal{L}$ -functionals) are again only defined up to an overall constant depending on k , so that differences are intrinsically defined, and the analogue of the scaling property (0.8) holds.

As is well-known, the volume ratio of L^2 -balls can be expressed as a Gram determinant:

$$\frac{\text{vol } \mathcal{B}^2(\mu, \phi)}{\text{vol } \mathcal{B}^2(\nu, \psi)} = \det \left(\langle s_i, s_j \rangle_{L^2(\nu, \psi)} \right)_{i,j}, \quad (0.9)$$

where $S = (s_1, \dots, s_N)$ is an $L^2(\mu, \phi)$ -orthonormal basis of $H^0(L)$ and $N := h^0(L)$. If we let

$$\det S \in H^0(X^N, L^{\boxtimes N})$$

be the determinant section locally defined by

$$(\det S)(x_1, \dots, x_N) := \det(s_i(x_j))_{i,j}, \quad (0.10)$$

a simple row and column expansion of the determinant (see [Dei99], Lemma 5.27 on p. 103 and formula (5.35) on p. 107) shows that the induced L^2 -norm satisfies

$$\|\det S\|_{L^2(\nu, \psi)}^2 = N! \det \left(\langle s_i, s_j \rangle_{L^2(\nu, \psi)} \right)_{i,j}. \quad (0.11)$$

On the other hand its L^∞ -counterpart

$$\|\det S\|_{L^\infty(K, \psi)}$$

does not depend on the chosen $L^2(\mu, \phi)$ -orthonormal basis S either and is closely related to the *transfinite diameter* for compact subsets of $\mathbb{C}^n$, introduced by

Leja in 1959 as a several complex variable extension of the classical notion from potential theory.

Indeed let $(X, L) = (\mathbb{P}^n, \mathcal{O}(1))$, and let $\phi = \psi$ be the weight induced by the section cutting out the hyperplane at infinity. If μ denotes the Haar measure on the compact torus $\mathbb{T}^n \subset \mathbb{C}^n$, then the family S_k of all monomials of degree $\leq k$ defines an $L^2(\mu, k\phi)$ -orthonormal basis of $H^0(X, kL)$.

If $K \subset \mathbb{C}^n$ is a compact subset, then (the log of) Leja's transfinite diameter is defined as

$$\lim_{k \rightarrow \infty} \frac{1}{k^{n+1}} \log \|\det S_k\|_{L^\infty(K, k\phi)}$$

up to a multiplicative constant. The existence of the limit in this setting, which corresponds classically to the *unweighted* situation, was in fact only proved in 1975 by Zaharjuta [Zah75].

An important property of the Haar measure μ on $\mathbb{T}^n$ is the so-called *Bernstein-Markov property*, which says that the distortion between the L^2 and L^∞ -norms has sub-exponential growth. We extend this notion to weighted measures in Section 2, where we show that (μ, ϕ) has the Bernstein-Markov if μ is a positive volume form (Lemma 2.2). If the weighted measure (μ, ϕ) with support E has the Bernstein-Markov property, then in particular we have (cf. Lemma 2.5)

$$\lim_{k \rightarrow \infty} \mathcal{L}_k(\mu, \phi) - \mathcal{L}_k(E, \phi) = 0,$$

which means that we can replace $\mathcal{L}_k(E, \phi)$ by $\mathcal{L}_k(\mu, \phi)$ in Theorem A if (μ, ϕ) has the Bernstein-Markov property. As a consequence, we will prove:

Corollary A. Let (μ, ϕ) be a weighted measure with the Bernstein-Markov property, and let E be its support. For each k , let S_k be an $L^2(\mu, k\phi)$ -orthonormal basis of $H^0(kL)$. Then for every weighted subset (K, ψ) we have

$$\lim_{k \rightarrow \infty} \frac{1}{kN_k} \log \|\det S_k\|_{L^\infty(K, k\psi)} = \mathcal{E}_{\text{eq}}(E, \phi) - \mathcal{E}_{\text{eq}}(K, \psi)$$

where E is the support of μ .

This says in particular that the left-hand side limit exists, which extends Zaharjuta's result. In the $\mathbb{C}^n$ case, the existence of the limit in the weighted case was also independently obtained in [BL07b] using [Rum07]. Corollary A shows that

$$\exp(-\mathcal{E}_{\text{eq}}(K, \psi))$$

can be viewed as the transfinite diameter of (K, ψ) .

We give in Proposition 3.7 a recursion formula relating the Aubin-Mabuchi energy on X to that on a hypersurface Y . It shows that Corollary A contains in particular Rumely's Robin-type formula for the transfinite diameter in $\mathbb{C}^n$ [Rum07]. We also show how to recover DeMarco-Rumely's results [DMR06] in Section 5.3.

0.5. Applications to Arakelov geometry. In the last part of the paper, we give two further applications of Theorems A and B related to Arakelov geometry. As a consequence of Theorem A, we will first describe the asymptotic behaviour of the Ray-Singer analytic torsion of a smooth weight with arbitrary curvature, refining results of Bismut-Vasserot [BV89].

Our second application is a generalisation of Yuan's equidistribution theorem for points of small height [Yua06] to the case of a *big* line bundle (but at archimedean places only). Assume that X is a smooth projective variety defined over a number field, say $\mathbb{Q}$ for simplicity. Let L be a big line bundle on $X/\mathbb{Q}$. Denoting by $\mathbb{A}$ the adèles of $\mathbb{Q}$, $H^0(kL)_{\mathbb{Q}}$ embeds as a co-compact subgroup of

$$H^0(kL)_{\mathbb{A}} \subset H^0(kL)_{\mathbb{R}} \times \prod_p H^0(L)_{\mathbb{Q}_p}$$

which enables us to normalise the Haar measure $\text{vol}_k^{\mathbb{A}}$ on $H^0(kL)_{\mathbb{A}}$ by

$$\text{vol}_k^{\mathbb{A}} H^0(kL)_{\mathbb{A}} / H^0(kL)_{\mathbb{Q}} = 1.$$

Suppose given once and for all an arbitrary collection (ϕ_p) of continuous weights on $L_{\mathbb{C}_p}$ over $X(\mathbb{C}_p)$ for every prime p , for instance those induced by a model of X over $\mathbb{Z}$. The superscript $\mathbb{A}$ will be used to indicate that an object implicitly depends on (ϕ_p) .

If ϕ is a continuous weight on $L_{\mathbb{C}}$ over $X(\mathbb{C})$ we define the *adelic unit ball*

$$\mathcal{B}^{\mathbb{A}}(\phi) := H^0(L)_{\mathbb{A}} \cap \left(\mathcal{B}_{\mathbb{R}}^{\infty}(\phi) \times \prod_p \mathcal{B}_{\mathbb{Q}_p}^{\infty}(\phi_p) \right) \quad (0.12)$$

and we can then consider the corresponding adelic $\mathcal{L}$ -functionals

$$\mathcal{L}_k^{\mathbb{A}}(\phi) := \frac{1}{kN_k} \log \text{vol}_k^{\mathbb{A}} \mathcal{B}^{\mathbb{A}}(k\phi) \quad (0.13)$$

As opposed to the other $\mathcal{L}$ -functionals introduced so far, the adelic $\mathcal{L}$ -functionals $\mathcal{L}_k^{\mathbb{A}}$ are well-defined without any further normalisation issue.

We now introduce the *adelic energy at equilibrium* as

$$\mathcal{E}_{\text{eq}}^{\mathbb{A}}(\phi) := \limsup_{k \rightarrow \infty} \mathcal{L}_k^{\mathbb{A}}(\phi) \in [-\infty, +\infty].$$

The exponential of the right-hand side is called the *sectional capacity* in [RLV00], where it is proved that the limsup actually is a limit when L ample. Still assuming that L is merely big, Theorems A and B together will enable us to show (Lemma 6.4) that $\mathcal{E}_{\text{eq}}^{\mathbb{A}}(\cdot)$ is differentiable at any weight ϕ where it is finite, with derivative given by integration against the equilibrium measure $\mu_{\text{eq}}(X(\mathbb{C}), \phi)$.

On the other hand, the above data allows to define the *height* $h_{\phi}^{\mathbb{A}}(x)$ of any point $x \in X(\overline{\mathbb{Q}})$ (cf. 6.4). If $x_j \in X(\overline{\mathbb{Q}})$ is a *generic* sequence, that is a sequence converging to the generic point of X in the Zariski topology, then it is an easy consequence of the adelic Minkowski theorem (cf. Section 6.2) that their heights admit the asymptotic lower bound

$$\liminf_{j \rightarrow \infty} h_{\phi}^{\mathbb{A}}(x_j) \geq \mathcal{E}_{\text{eq}}^{\mathbb{A}}(\phi).$$

Following the original variational principle first used by Szpiro, Ullmo and Zhang [SUZ97], we will prove

Theorem C. Using the above notations, suppose that $x_j \in X(\overline{\mathbb{Q}})$ is a generic sequence such that

$$\lim_{j \rightarrow \infty} h_{\phi}(x_j) = \mathcal{E}_{\text{eq}}^{\mathbb{A}}(\phi) \in \mathbb{R}.$$

Then the Galois orbits of the x_j 's equidistribute on $X(\mathbb{C})$ as $j \rightarrow \infty$ towards the equilibrium measure $\mu_{\text{eq}}(X(\mathbb{C}), \phi)$.

0.6. Theorems A and B: the case of an ample line bundle. The case of an *ample* line bundle L already covers the classical $\mathbb{C}^n$ situation. It might thus be of interest to know that somewhat simpler proofs of Theorems A and B can be provided in that case.

Let us first sketch the proof of Theorem A, assuming that L is ample. As a first remark, it is clearly enough to prove Theorem A for a fixed choice of (K_2, ϕ_2) (then take differences) and we can thus assume that $K_2 = X$ and ϕ_2 is smooth and strictly psh, i.e. $dd^c \phi_2 > 0$.

Step 1: Assume also that $K_1 = X$ and ϕ_1 is smooth and strictly psh. If we fix a smooth positive volume form μ on X , it is a simple consequence of the mean value inequality that each (μ, ϕ_j) satisfies the Bernstein-Markov property, and Lemma 2.5 shows that we may replace L^∞ -balls by L^2 -balls to prove Theorem A in that case. From there on the argument is quite similar to Donaldson's proof of Proposition 2 in [Don05a]. Indeed on the one hand the differential of $\mathcal{E}_{\text{eq}}(X, \cdot)$ at a smooth psh weight ψ is given by integration against $\mu_{\text{eq}}(X, \psi)$, which is just the smooth probability measure $\text{vol}(L)^{-1}(dd^c \psi)^n$ here. On the other hand, Lemma 4.1 shows that the differential of $\mathcal{L}_k(\mu, \cdot)$ at a smooth weight ψ is given by integration against the Bergman measure $\beta(\mu, k\psi)$ defined in Section 2.1. Now the Bouche-Catlin-Tian-Zelditch theorem implies that

$$\lim_{k \rightarrow \infty} \beta(\mu, k\psi) = \mu_{\text{eq}}(X, \psi),$$

that is: the differential of $\mathcal{L}_k(\mu, \cdot)$ converges to that of $\mathcal{E}_{\text{eq}}(X, \cdot)$ pointwise on smooth psh weights. Theorem A thus follows by integrating this convergence along the line segment between our two smooth psh weights ϕ_1 and ϕ_2 .

Step 2: Suppose now that (K_1, ϕ_1) is an arbitrary weighted subset (K, ϕ) . We have the following obvious equality between L^∞ -balls:

$$\mathcal{B}^\infty(K, \phi) = \mathcal{B}^\infty(X, \phi_K) \tag{0.14}$$

where ϕ_K is the supremum of all psh weights ψ such that $\psi \leq \phi$ on K . By Demailly's regularisation theorem we can find two sequences $\phi_j^\pm$ of smooth strictly psh weights converging monotonically to ϕ_K a.e. on X from above and below respectively. Since $\mathcal{L}_k(X, \cdot)$ is clearly non-decreasing, the present general case follows from Step 1 applied to $\phi_j^\pm$ by continuity of the Aubin-Yau energy along monotonic sequences, itself a consequence of Bedford-Taylor's continuity theorems for mixed Monge-Ampère operators on bounded psh functions.

Let us now sketch an alternative argument to prove Theorem B when L is ample, say in case $K = X$. If ϕ_1, ϕ_2 are smooth weights, Theorem A (which is already proved) and Lemma 2.5 imply that

$$\lim_{k \rightarrow \infty} \mathcal{L}_k(\mu, \phi_1) - \mathcal{L}_k(\mu, \phi_2) = \mathcal{E}_{\text{eq}}(X, \phi_1) - \mathcal{E}_{\text{eq}}(X, \phi_2).$$

On the other hand Lemma 4.1 says that the differential of $\mathcal{L}_k(\mu, \cdot)$ at a smooth weight ψ is given by integration against the Bergman measure $\beta(\mu, k\psi)$. Now a recent result by the first named author (cf. Theorem 2.1) implies that

$$\lim_{k \rightarrow \infty} \beta(\mu, k\psi) = \mu_{\text{eq}}(X, \psi)$$

holds for smooth weights ϕ with *arbitrary* curvature. Integrating along the line segment between ϕ_1 and ϕ_2 yields

$$\mathcal{E}_{\text{eq}}(X, \phi_1) - \mathcal{E}_{\text{eq}}(X, \phi_2) = \int_{t=0}^1 dt \int_X (\phi_1 - \phi_2) \mu_{\text{eq}}(X, t\phi_1 + (1-t)\phi_2),$$

which is equivalent to Theorem B (for $K = X$).

0.7. Structure of the paper.

- Sections 1 and 2 contain preliminary results on Monge-Ampère operators and Bergman kernels asymptotics.
- Section 3 extends to our singular setting standard facts on the Aubin-Mabuchi energy functional, and contains the proof of Theorem B.
- Section 4 contains the proofs of Theorem A and Corollary A.
- Section 5 presents applications to the $\mathbb{C}^n$ setting.
- Finally Section 6 presents applications to Arakelov geometry, in particular the proof of Theorem C.

Acknowledgement. We would like to thank B. Berndtsson, F. Berteloot, A. Chambert-Loir, J.-P. Demailly, V. Guedj, C. Mourougane and A. Zeriahi for interesting discussions related to the contents of the present paper. We are especially grateful to N. Levenberg for pointing out a gap in the proof of Theorem B in a previous version of this paper.

1. MIXED MONGE-AMPÈRE OPERATORS AND EQUILIBRIUM WEIGHTS

The goal of this section is to collect some results on mixed Monge-Ampère operators that are required to study the Aubin-Mabuchi energy functional in the case of a *big* line bundle L .

The reader primarily interested in the case of an *ample* line bundle will realise that the results we mention are completely standard in that setting (cf. for instance [GZ05]), and proofs in the general case can be found in [BEGZ08]

1.1. Big bundles and minimal singularities. Recall that a line bundle L on a compact complex manifold X is said to be *pseudo-effective* (*psef* for short) iff it admits a *psh* weight. The line bundle L is said to be *big* iff its *volume*

$$\text{vol}(L) := \limsup_{k \rightarrow \infty} \frac{n!}{k^n} h^0(kL)$$

is positive. Here we write as usual by $h^0 := \dim H^0$, and the $\limsup$ is actually a limit as a consequence of Fujita's theorem. A theorem independently proved by Bonavero [Bon98] and Ji-Shiffmann [JS93] asserts that L is big iff it admits a *strictly psh* weight, i.e. a singular weight ϕ whose curvature current $dd^c \phi$ dominates a (smooth) positive $(1, 1)$ -form.

It follows from Demailly's regularisation theorem [Dem92] that ϕ can then be chosen to have analytic singularities, and in particular to be locally bounded on a Zariski open subset Ω of X . Finally note that X is Moishezon, i.e. bimeromorphic to a projective manifold, iff it admits a big line bundle.

Given two *psh* weights ϕ_1, ϕ_2 on L , one says that ϕ_1 is *more singular* than ϕ_2 if $\phi_1 \leq \phi_2 + O(1)$. As has been observed by Demailly, any pseudo-effective

line bundle L admits psh weights with minimal singularities in this sense. Indeed given a smooth weight ϕ on L the equilibrium weight

$$P_X \phi = \sup \{ \psi, \psi \text{ psh weight on } L, \psi \leq \phi \}$$

is automatically (usc and) psh, and it plainly has minimal singularities. We will at any rate come back to this construction in what follows.

Note that the difference between any two psh weights with minimal singularities is a bounded function by definition. When L is ample, the psh weights with minimal singularities are exactly the locally bounded psh weights, and in the general case the former appear to share many of the nice properties the latter exhibit in the setting of pluripotential theory.

When L is only big, there exists as we saw a strictly psh weight that is locally bounded on a Zariski open subset Ω of X . It follows that *every* psh weight with minimal singularities on L is locally bounded on this same Ω .

The following easy result is proved in [BEGZ08].

Proposition 1.1. *Let $\pi : Y \rightarrow X$ be a surjective holomorphic map. If ϕ is a psh weight with minimal singularities on L , then $\pi^* \phi$ is a psh weight with minimal singularities on $\pi^* L$.*

1.2. Mixed Monge-Ampère operators and comparison principle. As explained above, results in this section are standard when dealing with ample line bundles. Indeed, they all follow from Bedford-Taylor's local results for locally bounded psh weights. The proofs in the general situation where line bundles are merely big can be found in [BEGZ08].

Let L be a big line bundle. By what we saw above, we can choose a Zariski open subset Ω on which every psh weight with minimal singularities is locally bounded.

Now let $\phi_1, \dots, \phi_n$ be psh weights on L that are locally bounded on Ω . We can then define the Bedford-Taylor wedge product

$$dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n$$

as a positive measure on Ω . Recall that this is done by locally setting $dd^c u \wedge T := dd^c(uT)$ whenever u is a locally bounded psh function and T is a closed positive current (which thus has measure coefficients). It was proved by Bedford-Taylor that the resulting measure $dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n$ puts no mass on pluripolar subsets of Ω . The following result is proved in [BEGZ08].

Theorem 1.2. *Let $\phi_1, \dots, \phi_n$ (resp. $\psi_1, \dots, \psi_n$) be psh weights on L that are locally bounded on a Zariski open subset Ω . If ϕ_j is less singular than ψ_j for all j , then we have*

$$\int_{\Omega} dd^c \psi_1 \wedge \dots \wedge dd^c \psi_n \leq \int_{\Omega} dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n \leq \text{vol}(L).$$

Equality holds on the right-hand side when the ϕ_j 's have minimal singularities.

This says in particular that $dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n$ has finite total mass, and we can thus introduce:

Definition 1.3. If $\phi_1, \dots, \phi_n$ are psh weights on L that are locally bounded on a Zariski open subset, the non-pluripolar product

$$\langle dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n \rangle$$

is defined as the trivial extension to X of the positive measure $dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n$ on Ω . In particular, the Monge-Ampère measure of a psh weight ϕ locally bounded on a Zariski open subset Ω is defined by

$$\text{MA}(\phi) := \langle (dd^c \phi)^n \rangle.$$

We stress that such non-pluripolar products $\langle dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n \rangle$ put no mass on pluripolar subsets of X , and therefore do not depend on the choice of Ω . By Theorem 1.2, the total mass

$$\int_X \langle dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n \rangle$$

only depends on the singularity classes of the ϕ_j 's and is equal to $\text{vol}(L)$ when the ϕ_j 's have minimal singularities.

The non-pluripolar Monge-Ampère operator so defined satisfies the following generalised comparison principle, which will be a crucial ingredient in the proof of Theorem B.

Corollary 1.4. *Let ϕ_1 and ϕ_2 be two psh weights on L that are locally bounded on a Zariski open subset. If $\phi_1 \leq \phi_2 + O(1)$, then we have*

$$\int_{\{\phi_2 < \phi_1\}} \text{MA}(\phi_1) \leq \int_{\{\phi_2 < \phi_1\}} \text{MA}(\phi_2).$$

Proof. It is an important result of Bedford-Taylor that $u \mapsto (dd^c u)^n$ is local in the plurifine topology for locally bounded psh functions u . By definition of the non-pluripolar Monge-Ampère operator, it follows that $\phi \mapsto \text{MA}(\phi)$ defined above is also local in the plurifine topology. Now let $\varepsilon > 0$. The psh weight with minimal singularities $\max(\phi_2, \phi_1 - \varepsilon)$ coincides with ϕ_2 on the plurifine open subset $\{\phi_2 > \phi_1 - \varepsilon\}$ and with $\phi_1 - \varepsilon$ on the plurifine open subset $\{\phi_2 < \phi_1 - \varepsilon\}$. It follows that

$$\begin{aligned} & \int_X \text{MA}(\max(\phi_2, \phi_1 - \varepsilon)) \\ & \geq \int_{\{\phi_2 > \phi_1 - \varepsilon\}} \text{MA}(\phi_2) + \int_{\{\phi_2 < \phi_1 - \varepsilon\}} \text{MA}(\phi_1) \end{aligned}$$

which is in turn

$$\geq \int_X \text{MA}(\phi_2) - \int_{\{\phi_2 < \phi_1 - \varepsilon\}} \text{MA}(\phi_2) + \int_{\{\phi_2 < \phi_1 - \varepsilon\}} \text{MA}(\phi_1).$$

On the other hand Theorem 1.2 yields

$$\int_X \text{MA}(\phi_2) = \int_X \text{MA}(\max(\phi_2, \phi_1 - \varepsilon))$$

since $\phi_1 \leq \phi_2 + O(1)$ implies

$$\max(\phi_2, \phi_1 - \varepsilon) = \phi_2 + O(1),$$

and the result now follows by monotone convergence by letting $\varepsilon \rightarrow 0$. $\square$

We infer the following domination principle (cf. [BEGZ08]):

Corollary 1.5. *Let ϕ_1 and ϕ_2 be two psh weights on L and suppose that ϕ_2 has minimal singularities. If $\phi_1 \leq \phi_2$ holds a.e. wrt $\text{MA}(\phi_1)$, then $\phi_1 \leq \phi_2$ everywhere on X .*

The following continuity result is proved in [BEGZ08].

Theorem 1.6. *Let ψ_0 be a fixed psh weight with minimal singularities on L . Then the measure-valued operators*

$$(\phi_1, \dots, \phi_n) \mapsto \langle dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n \rangle$$

and

$$(\phi_0, \dots, \phi_n) \mapsto (\phi_0 - \psi_0) \langle dd^c \phi_1 \wedge \dots \wedge dd^c \phi_n \rangle$$

are continuous along convergent sequence $\phi_j^{(k)} \rightarrow \phi_j$ of psh weights with minimal singularities in the following three cases:

- $\phi_j^{(k)}$ decreases pointwise to ϕ_j .
- $\phi_j^{(k)}$ increases to ϕ_j a.e. wrt Lebesgue measure.
- $\phi_j^{(k)}$ converges to ϕ_j uniformly on X .

For the first operator considered, this is in fact straightforward: convergence holds locally on the Zariski open subset Ω where weights are locally bounded by Bedford-Taylor's results, and it extends across the boundary of Ω because the total mass is constant by Theorem 1.2. The case of the second operator then follows quite easily.

The following integration-by-part formula is more difficult to establish. Its proof, given in [BEGZ08], is an elaboration of the Skoda-El Mir extension theorem.

Theorem 1.7. *Let u and v be two bounded functions on X , each being a differences of quasi-psh functions that are locally bounded on a given Zariski open subset Ω . Let also Θ be a closed positive current of bidimension $(1, 1)$ on X . Then we have*

$$\int_{\Omega} u dd^c v \wedge \Theta = \int_{\Omega} v dd^c u \wedge \Theta = - \int_{\Omega} dv \wedge d^c u \wedge \Theta.$$

1.3. Equilibrium weights. Let X be a compact complex manifold and L be a big line bundle. Given a weighted subset (K, ϕ) , we set

$$\phi_K = \sup \{ \psi, \psi \text{ psh weight on } L, \psi \leq \phi \text{ on } K \}, \quad (1.1)$$

so that the definition (0.3) of the equilibrium weight $P_K \phi$ can be reformulated as

$$P_K \phi = \phi_K^*.$$

In case $K = X$ the inequality $\phi_X \leq \phi$ on X implies $P_X \phi \leq \phi$ by continuity of ϕ , and this means that $P_X \phi = \phi_X$ is already upper semi-continuous in that case. This property however fails for more general weighted subsets. Extending the classical terminology, a weighted subset (K, ϕ) will be called *regular* if ϕ_K is usc, i.e if $P_K \phi \leq \phi$ holds on K .

By Choquet's lemma (cf. [KliBook] p. 38) there exists an increasing sequence of psh weights ψ_j such that $\psi_j \leq \phi$ on K and $\lim_{j \rightarrow \infty} \psi_j = P_K \phi$ a.e. on X wrt Lebesgue measure, and we can furthermore assume that the ψ_j have minimal singularities by replacing them by $\max(\psi_j, \tau)$ where τ is a psh weight with minimal singularities such that $\tau \leq \phi$ on K .

The following 'tautological maximum principle' is a mere reformulation of the definition of ϕ_K .

Proposition 1.8. (*Maximum principle*) *Let (K, ϕ) be weighted subset. Then for every psh weight ψ on L we have*

$$\sup_K (\psi - \phi) = \sup_X (\psi - \phi_K)$$

In particular,

$$\|s\|_{L^\infty(K, k\phi)} = \|s\|_{L^\infty(X, k\phi_K)}$$

for every section $s \in H^0(kL)$.

Note however that this fails in general with $P_K \phi$ in place of K when (K, ϕ) is not regular. As an immediate consequence of Proposition 1.1, we see that equilibrium weights behave well under pull-back, that is:

Proposition 1.9. *Let $\pi : Y \rightarrow X$ be a surjective morphism between two compact complex manifolds of same dimension n , and let L be a big line bundle on X (so that π^*L is also big). Let (K, ϕ) be a weighted subset of (X, L) , and consider the induced weighted subset $(\pi^{-1}K, \pi^*\phi)$ of (Y, π^*L) . Then their respective equilibrium weights are related by*

$$P_{\pi^{-1}K} \pi^* \phi = \pi^* P_K \phi.$$

Recall from (0.4) that the equilibrium measure of (K, ϕ) is defined by

$$\mu_{\text{eq}}(K, \phi) := \text{vol}(L)^{-1} \text{MA}(P_K \phi).$$

It is a probability measure by Theorem 1.2.

Proposition 1.10. *If (K, ϕ) is a weighted subset, then $\mu_{\text{eq}}(K, \phi)$ is concentrated on K and we have $P_K \phi = \phi$ on K a.e. with respect to this measure.*

The technique of proof is pretty standard (see e.g [DemPP], p.17), but we provide details since this result plays a crucial role in the proof of Theorem B.

Proof. Let ψ_j be a non-decreasing sequence of psh weights with minimal singularities converging to $P_K \phi$ a.e. wrt Lebesgue measure. Let B be an open ball contained in $\Omega - K$. Since ψ_j is bounded on B , by Bedford-Taylor we can find a bounded psh function τ_j on B such that $(dd^c \tau_j)^n = 0$ and which coincides with ψ_j on the boundary of B . Since τ_j can be written as a Perron envelope, it follows that $\tau_j \geq \psi_j$ and $\tau_{j+1} \geq \tau_j$ on B . Now let $\tilde{\psi}_j$ be the psh weight which coincides with ψ_j outside B and with τ_j on B . We then have $\tilde{\psi}_j = \psi_j \leq \phi$ on K since the latter doesn't meet B , hence

$$\psi_j \leq \tilde{\psi}_j \leq \phi_K \leq P_K \phi$$

by definition of ϕ_K . We thus see that $P_K\phi$ is the increasing limit a.e. of the psh weights $\tilde{\psi}_j$. Since we have $\text{MA}(\tilde{\psi}_j) = 0$ on B , it follows that $\text{MA}(P_K\phi) = 0$ on B by continuity of Monge-Ampère along monotonic sequences, and we have thus proved that $\mu_{\text{eq}}(K, \phi)$ is concentrated on K .

Let us now prove that $\text{MA}(P_K\phi)$ is also concentrated on the closed subset $\{P_K\phi \geq \phi\}$. The argument is essentially the same, except that we need to be slightly more careful to guarantee that $\tilde{\psi}_j \leq \phi$ on B . Let thus $x_0 \in \Omega$ such that $P_K\phi(x_0) < \phi(x_0) - \varepsilon$ with $\varepsilon > 0$. If B is a small open ball centered at x_0 , we can identify weights on $L|_B$ with functions. If B is small enough we have $P_K\phi < \phi(x_0) - \varepsilon$ on B by upper semi-continuity of $P_K\phi$ and $\phi \geq \phi(x_0) - \varepsilon$ by continuity of ϕ . If τ_j denotes as above the bounded psh function on B such that of $(dd^c\tau_j)^n = 0$ and which coincides with ψ_j on the boundary of B , then $\psi_j \leq \phi(x_0) - \varepsilon$ on B implies $\tau_j \leq \phi(x_0) - \varepsilon$ on the boundary of B , hence

$$\tau_j \leq \phi(x_0) - \varepsilon \leq \phi$$

on B by pluri-subharmonicity of τ_j (since $\phi(x_0) - \varepsilon$ is a constant). We thus see that $\tilde{\psi}_j$ defined as above satisfies $\tilde{\psi}_j \leq \phi$ on K , and the same reasoning as above yields $\text{MA}(P_K\phi) = 0$ on B as desired.

We have thus proved that $P_K\phi \leq \phi$ holds a.e. wrt $\text{MA}(P_K\phi)$. Conversely, the set $\{P_K\phi > \phi_K\}$ is negligible by definition hence pluripolar (cf. [DemPP]). Since we have $P_K\phi \geq \phi_K$ and $\text{MA}(P_K\phi)$ puts no mass on pluripolar subsets, it follows that $P_K\phi = \phi_K \leq \phi$ a.e. on K with respect to this measure, and the proof of the second point is complete. $\square$

We now quote from [Ber07b] the following description of $\mu_{\text{eq}}(X, \phi)$ for a C^2 weight ϕ on X since it implicitly plays a key role in the present paper (cf. the proof of Theorem 2.1):

Theorem 1.11. *If ϕ is a C^2 weight on L the equilibrium measure $\mu_{\text{eq}}(X, \phi)$ is absolutely continuous with respect to Lebesgue measure. In fact we have $dd^c\phi \geq 0$ pointwise on the compact subset $E := \{P_X\phi = \phi\}$, and*

$$\mu_{\text{eq}}(X, \phi) = \text{vol}(L)^{-1} \mathbf{1}_E (dd^c\phi)^n.$$

This result is in fact a simple consequence of the local $C^{1,1}$ regularity of $P_X\phi$ on a Zariski open subset, which is the technical core of [Ber07b].

1.4. Approximation by pluri-subharmonic envelopes of smooth weights.

Let K be a given compact non-pluripolar subset of X . We first record the following straightforward properties of the projection operator P_K .

Lemma 1.12. *The projection operator P_K is non-decreasing, concave and continuous along decreasing sequences of continuous weights on $L|_K$. It is also 1-Lipschitz continuous:*

$$\sup_X |P_K\phi_1 - P_K\phi_2| \leq \sup_K |\phi_1 - \phi_2|$$

for any two continuous weights ϕ_1, ϕ_2 on $L|_K$.

The following approximation result will allow us to reduce the proof of Theorem A to the case of smooth weights.

Proposition 1.13. *Let L be a big line bundle.*

- *Let ψ be a psh weight on L . Then there exists a decreasing sequence of smooth weights ϕ_j on L such that $\lim_{j \rightarrow \infty} P_X \phi_j = \psi$ pointwise on X .*
- *Let (K, ϕ) be a weighted subset. Then there exists an increasing sequence ϕ_j of smooth weights on L such that $\lim_{j \rightarrow \infty} P_X \phi_j = \phi_K$ almost everywhere wrt Lebesgue measure.*

Proof. Since ψ is in particular upper semi-continuous, we can find a decreasing sequence ϕ_j of smooth weights such that $\phi_j \rightarrow \psi$ pointwise on X . Since $\psi \leq \phi_j$ is psh, we infer $\psi \leq P_X \phi_j \leq \phi_j$, and it follows that $P_X \phi_j$ also decreases pointwise to ψ , which proves the first point.

Let us now prove the second point. We first claim that

$$\phi_K = \sup\{P_X \tau, \tau \text{ continuous weight on } L, P_X \tau \leq \phi \text{ on } K\}.$$

Indeed let ψ be psh weights such that $\psi \leq \phi$ on K and let $\varepsilon > 0$. By the first part of the proof, there exists a decreasing sequence ϕ_j of smooth weights such that $P_X \phi_j$ decreases pointwise to $\psi - \varepsilon$ as $j \rightarrow \infty$. By Dini's lemma, it follows that the usc function $P_X \phi_j - \phi$ is ≤ 0 on the compact set K for $j \gg 1$ large enough, and we thus get

$$\psi - \varepsilon \leq P_X \phi_j \leq \phi_K$$

for j large enough, hence the claim. Since the family of psh weights $P_X \tau$ as above is clearly stable by max, Choquet's lemma thus shows that there exists an increasing sequence τ_j of continuous weights such that $\tau_j \leq \phi$ on K and $P_X \tau_j \rightarrow \phi_K$ a.e. To conclude the proof we simply take an increasing sequence of smooth weight ϕ_j such that

$$\tau_j - 1/j \leq \phi_j \leq \tau_j.$$

□

Remark 1.14. When L is ample, it is not difficult to show that the smooth weights ϕ_j in both points of Proposition 1.13 can furthermore be taken *strictly psh*, and in particular $P_X \phi_j = \phi_j$. This shows in particular that

$$\phi_K = \sup\{\psi, \psi \text{ continuous psh weight on } L, \psi \leq \phi \text{ on } K\},$$

which is thus always lower semi-continuous in that case. It follows that (K, ϕ) is regular iff ϕ_K is continuous when L is ample, which corresponds to the classical definition (cf. [KliBook]).

2. THE BERGMAN DISTORTION FUNCTION AND THE BERNSTEIN-MARKOV PROPERTY

2.1. Bergman kernels. Let (μ, ϕ) be a weighted measure, and let E be the support of μ , which is non-pluripolar by our standing assumptions. The *Bergman distortion function* $\rho(\mu, \phi)$ is defined at a point $x \in E$ as the squared operator norm of the evaluation operator

$$\text{ev}_x : H^0(L) \rightarrow L_x,$$

in other words

$$\rho(\mu, \phi)(x) = \sup_{s \in H^0(L) - \{0\}} \frac{|s(x)|_\phi^2}{\|s\|_{L^2(\mu, \phi)}^2}. \quad (2.1)$$

Since μ is a probability measure we have

$$\|s\|_{L^2(\mu, \phi)} \leq \|s\|_{L^\infty(E, \phi)},$$

which shows that

$$\sup_E \rho(\mu, \phi)^{1/2}$$

is exactly the distortion between the $L^2(\mu, \phi)$ and $L^\infty(E, \phi)$ -norms on $H^0(L)$.

If $S = (s_1, \dots, s_N)$ denotes an $L^2(\mu, \phi)$ -orthonormal basis of $H^0(L)$, then it is well-known and easy to see that

$$\rho(\mu, \phi) = \sum_{j=1}^N |s_j|_\phi^2.$$

The *Bergman measure* associated to (μ, ϕ) is now defined as

$$\beta(\mu, \phi) := N^{-1} \rho(\mu, \phi) \mu. \quad (2.2)$$

Note that it is a *probability* measure since we have

$$\int_X \rho(\mu, \phi) \mu = \sum_j \|s_j\|_{L^2(\mu, \phi)}^2 = N.$$

If we now replace ϕ by $k\phi$, then the relation

$$\sup_K \rho(\mu, k\phi) \geq N_k$$

shows that the distortion between the $L^2(\mu, k\phi)$ and $L^\infty(E, k\phi)$ -norms on $H^0(kL)$ grows at least like $k^{n/2}$ as $k \rightarrow \infty$.

Assume now that μ is a smooth positive volume form on X and that ϕ is smooth, so that $E = X$ in particular. When ϕ has strictly positive curvature, the celebrated Bouche-Catlin-Tian-Zelditch theorem ([Bo90, Cat99, Tia90, Zel98]) asserts that $\beta(\mu, k\phi)$ admits a full asymptotic expansion in the space of smooth volume forms, with the probability measure $\mu_{\text{eq}}(X, \phi)$ as the dominant term.

As was shown by the first named author (in [Ber07a] for the $\mathbb{P}^n$ case and in [Ber07b] for the general case), part of this result still holds when the positive curvature assumption on ϕ is dropped.

Theorem 2.1. *Let L be a big line bundle, μ be a smooth positive volume form on X and ϕ be a C^2 weight on L . Then we have*

- $\sup_X \rho(\mu, \phi) = O(k^n)$.
- $\lim_{k \rightarrow \infty} \beta(\mu, k\phi) = \mu_{\text{eq}}(X, \phi)$ in the weak topology of measures.

Since this result plays a crucial role in what follows, we will sketch its proof for the convenience of the reader, and refer to [Ber07b] for the complete proof - a slightly more involved one in fact since Fujita's theorem is not used there but rather given a direct proof by analytic means.

Proof. By an elementary argument locally replacing ϕ by its second order Taylor expansion and using the mean value inequality, one shows that $\sup_X \rho(\mu, k\phi) = O(k^n)$, i.e. the first assertion, and

$$\limsup_{k \rightarrow \infty} N_k^{-1} \rho(\mu, k\phi) \mu \leq \text{vol}(L)^{-1} (dd^c \phi)^n$$

pointwise on the set where $dd^c \phi \geq 0$.

We now sketch the proof of the second point. Since we are dealing with probability measures, it is enough to show by weak compactness that if ν is a given accumulation point of the sequence of measures $\beta(\mu, k\phi)$, then necessarily $\nu \leq \mu_{\text{eq}}(\mu, \phi)$.

Now set $E := \{P_X \phi = \phi\}$, so that $dd^c \phi \geq 0$ on E and

$$\mu_{\text{eq}}(X, \phi) = \text{vol}(L)^{-1} \mathbf{1}_E (dd^c \phi)^n$$

by Theorem 1.11 recalled above from [Ber07b]. Since we automatically have

$$\rho(\mu, k\phi) \leq \exp(k(P_X \phi - \phi)) \sup_X \rho(\mu, k\phi)$$

by Proposition 1.8, the first assertion shows that $N_k^{-1} \rho(\mu, k\phi)$ tends to 0 (exponentially fast) pointwise on $X - E$.

Putting all this together yields

$$\limsup_{k \rightarrow \infty} N_k^{-1} \rho(\mu, k\phi) \mu \leq \mu_{\text{eq}}(X, \phi)$$

a.e. on X , and Lebesgue's dominated convergence finally implies that

$$\nu \leq \mu_{\text{eq}}(X, \phi)$$

for any accumulation point ν as desired. $\square$

2.2. The Bernstein-Markov property. Let μ be a positive volume form on X . By the first part of Theorem 2.1, if ϕ is a C^2 weight on L then the distortion

$$\sup_X \rho(\mu, k\phi)^{1/2}$$

between the $L^2(\mu, k\phi)$ and $L^\infty(X, k\phi)$ -norms on $H^0(kL)$ grows precisely like $k^{n/2}$, which is the minimal possible growth.

This fact is no longer true if we drop the smoothness assumption on ϕ . One can for instance show that the distortion is $O(k^{n/\alpha})$ when ϕ is of class C^α with $0 < \alpha < 2$, and this estimate is optimal. For general C^0 weights we have the following elementary fact:

Lemma 2.2. *Let μ be a smooth positive volume form. If ϕ is a C^0 weight on L , then the distortion has at most sub-exponential growth, i.e. for every $\varepsilon > 0$ we have*

$$\sup_X \rho(\mu, k\phi)^{1/2} = O(e^{\varepsilon k}).$$

Proof. Given $\varepsilon > 0$ we can cover X by a finite number of small balls B_j on which L is trivialised and ϕ is ε -close to its value ϕ_j at the centre of the ball. We can also assume that X is still covered by smaller balls B'_j relatively compact in B_j .

A section $s \in H^0(kL)$ is identified with a holomorphic section on B_j , and the desired inequality

$$|s(x)|^2 e^{-2k\phi} \leq C e^{2\epsilon k} \int_{B_j} |s|^2 e^{-2k\phi} d\mu$$

for $x \in B'_j$ is thus an immediate consequence of the mean value inequality applied to the psh function $|s|^2 e^{-2k\phi_j}$ on B_j . $\square$

We introduce the following extension of standard terminology (see [BL07a]):

Definition 2.3. A weighted measure (μ, ϕ) is said to have the *Bernstein-Markov property* if the distortion between the $L^2(\mu, k\phi)$ and $L^\infty(E, k\phi)$ -norms on $H^0(kL)$ has sub-exponential growth (E denoting the support of μ).

The following result is shown in [BBWN08], generalising results of Siciak [Sic88].

Theorem 2.4. *Let (E, ϕ) be a weighted subset and let μ be a probability measure supported on E . Assume that:*

- (E, ϕ) is regular, i.e. $P_E \phi \leq \phi$ holds on E .
- μ is determining for (E, ϕ) , i.e. for every psh weight ψ on L , $\psi \leq \phi$ a.e. wrt μ already implies $\psi \leq \phi$ on E .

Then (μ, ϕ) has the Bernstein-Markov property.

This somewhat technical looking criterion actually shows that a host of reasonable measures satisfy the Bernstein-Markov property. If E is for instance either (the closure of) a smoothly bounded domain or a totally real submanifold of X , then (E, ϕ) is regular for any continuous weight ϕ on $L|_E$. By the domination principle, if μ is a probability measure with positive C^0 density on a smoothly bounded domain or if μ is the equilibrium measure of a totally real submanifold E , we thus get that (μ, ϕ) satisfies the Bernstein-Markov property.

In the present article, we shall actually only use the following two special cases of Theorem 2.4: either μ and ϕ are both smooth (in which case the Bernstein-Markov property is a trivial consequence of the mean value inequality, as already noticed), or μ is the Haar measure on the unit compact torus $\mathbb{T}^n \subset (\mathbb{C}^*)^n \subset \mathbb{P}^n$, in which case the result is standard.

The next lemma will allow us to replace L^∞ -balls by L^2 -balls whenever convenient.

Lemma 2.5. *Let (μ, ϕ) be a weighted measure and let E be the support of μ . Then we have*

$$0 \leq \mathcal{L}_k(\mu, \phi) - \mathcal{L}_k(E, \phi) \leq \frac{1}{2k} \log \sup_E \rho(\mu, k\phi).$$

In particular if (μ, ϕ) is Bernstein-Markov then

$$\lim_{k \rightarrow \infty} \mathcal{L}_k(\mu, \phi) - \mathcal{L}_k(E, \phi) = 0.$$

Proof. If we set

$$C_k := \frac{1}{2k} \log \sup_E \rho(\mu, k\phi)$$

then we have

$$\|s\|_{L^2(\mu, k\phi)} \leq \|s\|_{L^\infty(\mu, k\phi)} \leq e^{kC_k} \|s\|_{L^2(\mu, k\phi)}$$

for all k and all sections $s \in H^0(kL)$. Since the volume form vol_k is homogeneous of degree $2N_k$ on $H^0(kL)$ we get

$$0 \leq \log \frac{\text{vol}_k \mathcal{B}^2(\mu, k\phi)}{\text{vol}_k \mathcal{B}^\infty(E, k\phi)} \leq 2kN_k C_k$$

and the result follows by definition (0.7) and its L^2 -analogue. $\square$

3. THE AUBIN-MABUCHI ENERGY FUNCTIONAL

In this section L denotes a big line bundle on X . We have chosen to use the language of weights in this section since the rest of the article is naturally expressed in this language, but it is of course immediate to extend the results of this section (and Theorem B in particular) to the more general case of θ -psh functions, where θ is a closed smooth $(1, 1)$ -form with big cohomology class.

3.1. The energy functional on psh weights. Let us first fix a psh weight ψ_0 with minimal singularities. As explained in the introduction, we define the Aubin-Mabuchi functional $\mathcal{E}$ on psh weights with minimal singularities by the formula

$$\mathcal{E}(\phi) = \frac{1}{n+1} \sum_{j=0}^n \int_X (\phi - \psi) \langle (dd^c \phi)^j \wedge (dd^c \psi_0^{n-j}) \rangle$$

This normalises $\mathcal{E}$ by the condition $\mathcal{E}(\psi_0) = 0$.

As in Section 1.2, the brackets denote non-pluripolar products. Concretely this means that the integrals are only extended over a Zariski open subset Ω of X on which all psh weights are locally bounded, so that Bedford-Taylor wedge products are well-defined on Ω .

We now verify that $\mathcal{E}$ remains a primitive of the Monge-Ampère operator in our singular setting,

Proposition 3.1. *For any two psh weights ϕ_1, ϕ_2 with minimal singularities we have*

$$\frac{d}{dt}_{t=0+} \mathcal{E}((1-t)\phi_1 + t\phi_2) = \int_X (\phi_2 - \phi_1) \text{MA}(\phi_1)$$

Proof. The function $u := \phi_2 - \phi_1$ is bounded. We compute

$$\begin{aligned} (n+1)\mathcal{E}((1-t)\phi_1 + t\phi_2) &= \int_\Omega (\phi_1 - \psi_0 + tu) \sum_{j=0}^n ((1-t)dd^c \phi_1 + tdd^c \phi_2)^j \wedge (dd^c \psi_0)^{n-j} \\ &= \int_\Omega (\phi_1 - \psi_0) \sum_{j=0}^n (dd^c \phi_1)^j \wedge (dd^c \psi_0)^{n-j} \\ &\quad + t \int_\Omega u \sum_{j=0}^n (dd^c \phi_1)^j \wedge (dd^c \psi_0)^{n-j} \end{aligned}$$

$$+t \int_{\Omega} (\phi_1 - \psi_0) \sum_{j=1}^n j (dd^c \phi_1)^{j-1} \wedge dd^c u \wedge (dd^c \psi_0)^{n-j} + O(t^2).$$

By integration-by-parts (Theorem 1.7) we have

$$\begin{aligned} \int_{\Omega} (\phi_1 - \psi_0) \sum_{j=1}^n j (dd^c \phi_1)^{j-1} \wedge dd^c u \wedge (dd^c \psi_0)^{n-j} &= \int_{\Omega} u dd^c (\phi_1 - \psi_0) \sum_{j=1}^n j (dd^c \phi_1)^{j-1} \wedge (dd^c \psi_0)^{n-j} \\ &= \int_{\Omega} u \sum_{j=1}^n j (dd^c \phi_1)^j \wedge (dd^c \psi_0)^{n-j} - \int_X u \sum_{j=0}^{n-1} (j+1) (dd^c \phi_1)^j \wedge (dd^c \psi_0)^{n-j}. \end{aligned}$$

Now

$$\begin{aligned} \sum_{j=0}^n (dd^c \phi_1)^j \wedge (dd^c \psi_0)^{n-j} + \sum_{j=1}^n j (dd^c \phi_1)^j \wedge (dd^c \psi_0)^{n-j} - \sum_{j=0}^{n-1} (j+1) (dd^c \phi_1)^j \wedge (dd^c \psi_0)^{n-j} \\ = (dd^c \phi_1)^n + n (dd^c \phi_1)^n \end{aligned}$$

It follows that

$$\mathcal{E}((1-t)\phi_1 + t\phi_2) = \mathcal{E}(\phi_1) + t \int_{\Omega} u (dd^c \phi_1)^n + O(t^2)$$

as desired. $\square$

As a consequence, we see that (0.2) always holds, that is:

Corollary 3.2. *For any two psh weights with minimal singularities ϕ, ψ we have*

$$\mathcal{E}(\phi) - \mathcal{E}(\psi) = \frac{1}{n+1} \sum_{j=0}^n \int_X (\phi - \psi) \langle (dd^c \phi)^j \wedge (dd^c \psi)^{n-j} \rangle.$$

Proof. We fix ψ and temporarily denote by $\mathcal{F}(\phi)$ the right-hand side expression. We can then apply Proposition 3.1 with ψ in place of ϕ_0 to get

$$\frac{d}{dt} \mathcal{F}((1-t)\phi + t\psi) = \int_X (\psi - \phi) \text{MA}((1-t)\phi + t\psi) = \frac{d}{dt} \mathcal{E}((1-t)\phi + t\psi),$$

and the result follows since $\mathcal{F}(\cdot)$ and $\mathcal{E}(\cdot) - \mathcal{E}(\psi)$ both vanish at ψ . $\square$

3.2. General properties of the energy. Theorem 1.6 implies the following continuity properties of the energy:

Proposition 3.3. *The map $\phi \mapsto \mathcal{E}(\phi)$ is continuous along converging sequences $\phi_j \rightarrow \phi$ of psh weights with minimal singularities in the following three cases.*

- ϕ_j decreases to ϕ pointwise.
- ϕ_j increases to ϕ a.e. for the Lebesgue measure.
- ϕ_j converges to ϕ uniformly on X .

Proposition 3.4. *The map $\phi \mapsto \mathcal{E}(\phi)$ is non-decreasing and concave on psh weights with minimal singularities.*

Proof. The first point follows from Corollary 3.2. To prove concavity, let ϕ_1, ϕ_2 be two psh weights with minimal singularities and set

$$g(t) := \mathcal{E}(t\phi_1 + (1-t)\phi_2).$$

By Proposition 3.1, we have

$$g'(t) = \int_X u \operatorname{MA}((1-t)\phi_1 + t\phi_2)$$

with $u := \phi_2 - \phi_1$. Computing the second derivative yields

$$\begin{aligned} g''(t) &= n \int_{\Omega} u dd^c u \wedge ((1-t)dd^c \phi_1 + tdd^c \phi_2)^{n-1} \\ &= -n \int_{\Omega} du \wedge d^c u \wedge ((1-t)dd^c \phi_1 + tdd^c \phi_2)^{n-1} \leq 0 \end{aligned}$$

by Theorem 1.7 again, and the proof is complete. $\square$

Remark 3.5. More generally one can consider variations along a 1-parameter family ϕ_t (with t in the unit-disc Δ in $\mathbb{C}$) of weights on L with minimal singularities. Under suitable regularity assumptions on $(t, x) \mapsto \phi_t(x)$ a simple modification of the previous proof yields

$$dd_t^c \mathcal{E}(\phi_t) = \int_{x \in X} (dd_{(x,t)}^c \phi_t(x))^{n+1}, \quad (3.1)$$

In the smooth case at least, this formula is well-known in Kähler geometry. When L is ample the operator that maps a curve ϕ_t of smooth strictly psh weights to the Monge-Ampère measure $(dd_{(x,t)}^c \phi_t(x))^{n+1}$ may be identified with the geodesic curvature of the curve $dd_x^c \phi_t$ in the space of all Kähler metrics $\mathcal{K}(X, L)$ on X lying in the first Chern class $c_1(L)$. The geodesic curvature is defined with respect to the Riemannian metric on $\mathcal{K}(X, L)$ naturally defined at ϕ by taking L^2 norms of tangent vectors with respect to the volume form $(dd_x^c \phi)^n$ [Che00]. Formula (3.1) thus shows that $\mathcal{E}$ is *affine* along geodesic segments in $\mathcal{K}(X, L)$.

It is also interesting to note that

$$\frac{k^{n+1}}{(n+1)!} \int_X (dd_{(x,t)}^c \phi)^{n+1}$$

is the leading term of the $(1, 1)$ -part of the pushed-forward form

$$\int_X \operatorname{ch}_{X \times \Delta}(kL, k\phi) \operatorname{td}(T_X, \omega),$$

which coincides with the curvature of the Quillen metric on $\det H^\bullet(kL)$ by the main result of [BGS88] (see also [SouBook], Theorem 4 p.132).

Remark 3.6. As a consequence of Proposition 3.4, we may extend as in [BEGZ08] the definition of $\mathcal{E}(\phi)$ to an *arbitrary* psh weight on L as follows:

$$\mathcal{E}(\phi) = \inf_{\psi \geq \phi} \mathcal{E}(\psi) \in [-\infty, +\infty[$$

for ψ ranging over all psh weights with minimal singularities such that $\psi \geq \phi$. It is straightforward to see that $\phi \mapsto \mathcal{E}(\phi)$ so defined remains non-decreasing and concave on all psh weights. It is shown in [BEGZ08] that it is also upper

semi-continuous in the weak topology and that Corollary 3.2 remains true if both $\mathcal{E}(\phi)$ and $\mathcal{E}(\psi)$ are finite.

The following result relates the Aubin-Mabuchi energy $\mathcal{E}_X$ on X to the energy $\mathcal{E}_Y$ on a hypersurface Y of X . We assume here that L is ample and Y is smooth for simplicity.

Proposition 3.7. *Let L be an ample line bundle, and assume that Y is a smooth hypersurface of X cut out by a section $s \in H^0(X, L)$. If ϕ, ψ are locally bounded psh weights on L then we have*

$$\begin{aligned} \mathcal{E}_X(\phi) - \mathcal{E}_X(\psi) &= \mathcal{E}_Y(\phi|_Y) - \mathcal{E}_Y(\psi|_Y) \\ &+ \frac{1}{n+1} \left(\int_X \log |s|_\psi \text{MA}(\psi) - \int_X \log |s|_\phi \text{MA}(\phi) \right). \end{aligned}$$

Proof. Consider the following simple algebraic formula

$$(dd^c \phi)^n - (dd^c \psi)^n = dd^c((\phi - \psi) \sum_{j=0}^{n-1} (dd^c \phi)^j \wedge (dd^c \psi)^{n-1-j}), \quad (3.2)$$

From the point of view of Bott-Chern secondary characteristic classes, it may be interpreted as a double transgression formula (compare [Don85, SouBook]). At any rate, multiplying (3.2) by

$$u_\varepsilon := \log(|s|_\phi + \varepsilon)$$

and using integration by parts gives

$$\begin{aligned} \int_X u_\varepsilon (dd^c \phi)^n - u_\varepsilon (dd^c \psi)^n + (dd^c \phi) \wedge (\phi - \psi) \sum_{j=0}^{n-1} (dd^c \phi)^j \wedge (dd^c \psi)^{n-1-j} \\ = \int_X (\psi - \phi) dd^c(u_\varepsilon + \phi) \wedge \sum_{j=0}^{n-1} (dd^c \phi)^j \wedge (dd^c \psi)^{n-1-j}. \end{aligned}$$

Now $u_\varepsilon + \phi$ decreases to $\log |s|$, thus this converges to

$$(n+1) (\mathcal{E}_Y(\phi|_Y) - \mathcal{E}_Y(\psi|_Y))$$

by the the Lelong-Poincaré formula, and the result easily follows. $\square$

The following pull-back formula is straightforward using Proposition 1.1.

Proposition 3.8. *Let $\pi : Y \rightarrow X$ be a surjective morphism between compact complex manifolds of same dimension n and denote by e its (topological) degree. Let L be a big line bundle on X , and let ϕ, ψ be two psh weights with minimal singularities on L . Then we have*

$$\mathcal{E}_Y(\pi^* \phi) - \mathcal{E}_Y(\pi^* \psi) = e (\mathcal{E}_X(\phi) - \mathcal{E}_X(\psi)).$$

3.3. Proof of Theorem B. In this section we will prove Theorem B. Let thus K be a non-pluripolar compact subset of X . We first prove that

$$\phi \mapsto \mathcal{E}_{\text{eq}}(K, \phi) = \text{vol}(L)^{-1} \mathcal{E} \circ P_K(\phi)$$

is concave and continuous. Concavity is an immediate consequence of Proposition 3.4: for any weights ϕ_1, ϕ_2 on $L|_K$ we have

$$P_K((1-t)\phi_1 + t\phi_2) \geq (1-t)P_K\phi_1 + tP_K\phi_2$$

by concavity of P_K (Lemma 1.12) hence

$$\mathcal{E}(P_K((1-t)\phi_1 + t\phi_2)) \geq \mathcal{E}((1-t)P_K\phi_1 + tP_K\phi_2)$$

since $\mathcal{E}$ is non-decreasing

$$\geq (1-t)\mathcal{E}(P_K\phi_1) + t\mathcal{E}(P_K\phi_2)$$

since $\mathcal{E}$ is concave. Continuity of $\phi \mapsto \mathcal{E}_{\text{eq}}(K, \phi)$ follows from Lemma 1.12 and the third case of Proposition 3.3.

Given a continuous weight ϕ on $L|_K$ and $u \in C^0(K)$, the concave function $\mathcal{E} \circ P_K$ admits a directional derivative at ϕ in the direction u , and our goal is to show that it is given by

$$\frac{d}{dt}_{t=0_+} \mathcal{E} \circ P_K(\phi + tu) = \langle \lambda, u \rangle$$

where λ is the linear form on $C^0(K)$ defined by

$$\langle \lambda, u \rangle = \int_K u \text{MA}(P_K\phi).$$

Note that λ computes the directional derivatives of $\mathcal{E}$ at $P_K\phi$ according to Proposition 3.1.

The problem at hand is an instance of a differentiability property for the optimal value of a concave optimisation problem with parameter. Indeed since $\mathcal{E}$ is non-decreasing we have

$$\mathcal{E} \circ P_K(\phi) = \sup\{\mathcal{E}(\psi), \psi \text{ psh weight with minimal singularities } \psi \leq \phi \text{ on } K\}$$

by Choquet's lemma and continuity of the energy along non-decreasing sequences.

There has been a certain amount of work on differentiability of such optimal values in an abstract setting, but it seems that what we are trying to prove doesn't follow formally from such general results. On the other, the proof of Lemma 3.9, though pretty elementary, was inspired by more delicate considerations in [LM80].

The next lemma enables us to replace $\mathcal{E}$ by its linearisation λ at $P_K\phi$.

Lemma 3.9. *We have*

$$\frac{d}{dt}_{t=0_+} \mathcal{E} \circ P_K(\phi + tu) = \frac{d}{dt}_{t=0_+} \langle \lambda, P_K(\phi + tu) - P_K\phi \rangle.$$

Proof. Set for simplicity

$$a := \frac{d}{dt}_{t=0_+} \langle \lambda, P_K(\phi + tu) - P_K\phi \rangle,$$

which exists since $\lambda \circ P_K$ is concave. On the one hand, concavity of $\mathcal{E}$ yields

$$\mathcal{E} \circ P_K(\phi + tu) \leq \mathcal{E} \circ P_K(\phi) + \langle \lambda, P_K(\phi + tu) - P_K\phi \rangle,$$

hence

$$\frac{d}{dt}_{t=0+} \mathcal{E} \circ P_K(\phi + tu) \leq a.$$

On the other hand, given $\varepsilon > 0$ we can fix $\delta > 0$ small enough such that

$$\langle \lambda, P_K(\phi + \delta u) - P_K(\phi) \rangle \geq \delta a - \delta \varepsilon. \quad (3.3)$$

For $t > 0$ small enough we then have

$$\mathcal{E}((1-t)P_K\phi + tP_K(\phi + \delta u)) \geq \mathcal{E} \circ P_K(\phi) + t\langle \lambda, P_K(\phi + \delta u) - P_K\phi \rangle - t\delta\varepsilon$$

by Proposition 3.1

$$\geq \mathcal{E} \circ P_K(\phi) + t\delta a - 2t\delta\varepsilon$$

by (3.3). But since

$$P_K(\phi + t\delta u) \geq (1-t)P_K\phi + tP_K(\phi + \delta u)$$

by concavity of P_K , we finally infer that

$$\mathcal{E} \circ P_K(\phi + t\delta u) \geq \mathcal{E} \circ P_K(\phi) + t\delta a - 2t\delta\varepsilon$$

for all $t > 0$ small enough by monotonicity of $\mathcal{E}$. It follows that

$$\frac{d}{dt}_{t=0+} \mathcal{E} \circ P_K(\phi + tu) \geq a - 2\varepsilon$$

for each $\varepsilon > 0$, and the result follows. $\square$

We are now reduced to proving the linearised version of the problem, to wit

Lemma 3.10. *The super-differential at ϕ of the linearised problem is reduced to λ . In other words, we have*

$$\frac{d}{dt}_{t=0+} \langle \lambda, P_K(\phi + tu) - P_K\phi \rangle = \langle \lambda, u \rangle$$

for each $u \in C^0(K)$.

A crucial ingredient here is the following *orthogonality relation*

$$\langle \lambda, P_K\phi - \phi \rangle = 0,$$

which follows from Proposition 1.10. Since $P_K(\phi + u) \leq \phi + u$, this property implies

$$\langle \lambda, P_K(\phi + u) - P_K(\phi) \rangle \leq \langle \lambda, u \rangle,$$

which means that the linear form λ belongs to the super-differential at 0 of the continuous concave function

$$u \mapsto \langle \lambda, P_K(\phi + u) - P_K\phi \rangle.$$

At this point, we also see that Theorem B reduces to the differentiability part of the assertion, since the differential then has to coincide with λ .

Proof. We now prove Lemma 3.10. Our goal is to show that

$$\langle \lambda, P_K(\phi + tu) - P_K\phi - tu \rangle = o(t).$$

Since on the one hand

$$P_K(\phi + tu) \leq \phi + tu = P_K\phi + tu$$

λ -a.e. and on the other hand

$$\sup_X |P_K(\phi + tu) - P_K\phi - tu| = O(t)$$

by Lemma 1.12, it will be enough to show that

$$\lim_{t \rightarrow 0+} \int_{\{P_K(\phi + tu) < P_K\phi + tu\}} \lambda = 0. \quad (3.4)$$

We are going to show this by applying the comparison principle (Corollary 1.4).

We first claim that $u \in C^0(K)$ may be assumed to be the restriction to K of a *smooth* function on X . Indeed Theorem B admits the following integral reformulation

$$\mathcal{E} \circ P_K(\phi + u) - \mathcal{E} \circ P_K(\phi) = \int_{t=0}^1 \int_K u \operatorname{MA}(P_K(\phi + tu)) dt.$$

If we let u_j be a sequence of smooth functions on X converging uniformly to u on K , then it is $P_K(\phi + tu_j) \rightarrow P_K(\phi + tu)$ uniformly on X by Lemma 1.12. By Proposition 3.3, we deduce that

$$\lim_{j \rightarrow \infty} \mathcal{E} \circ P_K(\phi + u_j) = \mathcal{E} \circ P_K(\phi + u).$$

On the other hand for each t we have

$$\begin{aligned} & \int_K u_j \operatorname{MA}(P_K(\phi + tu_j)) - \int_K u \operatorname{MA}(P_K(\phi + tu)) \\ &= \int_K (u_j - u) \operatorname{MA}(P_K(\phi + tu_j)) + \int_K u (\operatorname{MA}(P_K(\phi + tu_j)) - \operatorname{MA}(P_K(\phi + tu))). \end{aligned}$$

The first term is bounded by $\operatorname{vol}(L) \sup_K |u_j - u|$ by Theorem 1.2, whereas the second one converges to 0 by Theorem 1.6. We thus see that

$$\lim_{j \rightarrow \infty} \int_K u_j \operatorname{MA}(P_K(\phi + tu_j)) = \int_K u \operatorname{MA}(P_K(\phi + tu))$$

for all t , and we infer

$$\lim_{j \rightarrow \infty} \int_{t=0}^1 \int_K u_j \operatorname{MA}(P_K(\phi + tu_j)) dt = \int_{t=0}^1 \int_K u \operatorname{MA}(P_K(\phi + tu)) dt$$

by dominated convergence, which shows our claim.

We now fix a strictly psh weight ϕ_+ with analytic singularities on L . Since u is smooth, it follows that $\phi_+ + \varepsilon u$ is psh for $\varepsilon > 0$ small enough. Upon scaling u , we may assume that $\varepsilon = 1$.

Since $P_K\phi = P_K(\phi + tu) + O(t)$ by Lemma 1.12, it follows in particular that

$$P_K(\phi) + t\phi_+ + tu$$

and

$$P_K(\phi + tu) + t\phi_+$$

have equivalent singularities, and the generalised comparison principle (Corollary 1.4) thus yields

$$\begin{aligned} & \int_{O_t} \text{MA} (P_K(\phi) + t(u + \phi_+)) \\ & \leq \int_{O_t} \text{MA} (P_K(\phi + tu) + t\phi_+) \end{aligned}$$

with

$$O_t := \{P_K(\phi + tu) < P_K(\phi) + tu\}.$$

Now we have

$$\text{MA} (P_K(\phi + tu) + t\phi_+) = \text{MA} (P_K(\phi + tu)) + \sum_{j=1}^n \binom{n}{j} t^j (dd^c P_K(\phi + tu))^{n-j} \wedge (dd^c \phi_+)^j \quad (3.5)$$

on the Zariski open subset where everyone is locally bounded by the binomial formula. Since $t(u + \phi_+)$ is psh by assumption, we have

$$\int_{O_t} \text{MA} (P_K \phi) \leq \int_{O_t} \text{MA} (P_K(\phi) + t(u + \phi_+)),$$

which is in turn

$$\leq \int_{O_t} \text{MA} (P_K(\phi + tu)) + O(t)$$

by (3.5) and Theorem 1.2. But $P_K \phi \leq \phi$ implies that

$$O_t \subset \{P_K(\phi + tu) < \phi + tu\},$$

and we infer that

$$\int_{O_t} \text{MA} (P_K(\phi + tu)) = 0$$

by Proposition 1.10 again. We thus conclude that

$$\int_{O_t} \text{MA} (P_K \phi) = O(t),$$

and the proof of Lemma 3.10 is thus complete. $\square$

We now show that the energy at equilibrium is $C^{1,1}$ in the following sense:

Proposition 3.11. *Let (K, ϕ) be a weighted subset and let $u \in C^0(K)$ be the restriction to K of a smooth function on X . Then $t \mapsto \mathcal{E}_{\text{eq}}(K, \phi + tu)$ is of class $C^{1,1}$ on $\mathbb{R}$.*

Proof. By Theorem B the derivative is equal (up to the factor $\text{vol}(L)^{-1}$) to

$$g(t) := \int_X u \text{MA} (P_K(\phi + tu)),$$

and we are to show that g is locally Lipschitz continuous on $\mathbb{R}$. But (3.2) above and integration by parts (Theorem 1.7) yield

$$g(t) - g(s) = \int_X (P_K(\phi + tu) - P_K(\phi + su)) dd^c u \wedge \Theta_t$$

with

$$\Theta_t := \sum_{j=1}^{n-1} \langle (dd^c P(\phi + tu))^j \wedge (dd^c P(\phi + su))^{n-j} \rangle.$$

But we have

$$\sup_X |P_K(\phi + tu) - P_K(\phi + su)| \leq C|t - s|$$

by Lemma 1.12, and on the other hand the total mass of the positive currents Θ_t have uniformly bounded mass by Theorem 1.2, and the result follows. $\square$

Example 3.12. Even for $K = X$ the energy is not twice differentiable in general. Indeed let $\chi, \theta \geq 0$ be smooth non-negative functions on $\mathbb{R}$ such that:

- χ is convex, $\chi(x) = 0$ iff $x \leq 1$, $\chi(x) = x$ for $x \geq 2$.
- $\theta(x) = 1$ for $x \leq \varepsilon$, $\theta(x) = 0$ for $x \geq 1$ and θ is strictly concave on $]\varepsilon, 1 - \varepsilon[$, strictly convex on $]1 - \varepsilon, 1[$ for $\varepsilon > 0$ small enough.

Let χ_t be the convex envelope from below of $\chi + t\theta$. For $t \geq 0$ we have $\chi_t = \chi + t\theta$. For $t \leq 0$ we have $\chi_t = -t\theta$ on

$$\{\chi_t = \chi + t\theta\} \cap \text{supp } \theta$$

and $\chi_t = \chi$ for x large enough (draw a picture!).

The smooth subharmonic function $\chi(\log |z|)$ on $\mathbb{C}$ coincides with $\log |z|$ near infinity, hence extends to a smooth psh weight ϕ on $\mathcal{O}(1)$. On the other hand $u(z) = \theta(\log |z|)$ vanishes near infinity, hence extends to a smooth function on $\mathbb{P}^1$. Radial symmetry ensures that

$$\mu_{\text{eq}}(\mathbb{P}^1, \phi + tu) = dd^c \chi_t(\log |z|)$$

on $\mathbb{C}$. The derivative $g(t)$ of $t \mapsto \mathcal{E}_{\text{eq}}(\mathbb{P}^1, \phi + tu)$ is thus given by $g(t) = 0$ for $t \geq 0$ and

$$g(t) = \int u dd^c \chi_t(\log |z|) = -t \int u dd^c u$$

for $t \leq 0$ by the above considerations. We thus get

$$g'(0_-) = \int du \wedge d^c u > 0 = g'(0_+),$$

which shows that g is not differentiable at 0.

4. VOLUME GROWTH AND TRANSFINITE DIAMETER

4.1. Proof of Theorem A. Let (K_1, ϕ_1) and (K_2, ϕ_2) be two weighted subset. Our goal is to prove that

$$\lim_{k \rightarrow \infty} \mathcal{L}_k(K_1, \phi_1) - \mathcal{L}_k(K_2, \phi_2) = \mathcal{E}_{\text{eq}}(K_1, \phi_1) - \mathcal{E}_{\text{eq}}(K_2, \phi_2). \quad (4.1)$$

If this formula holds for all (K_1, ϕ_1) and a *fixed* (K_2, ϕ_2) , then it also holds for *any* (K_2, ϕ_2) by taking differences. We can thus assume that $K_2 = X$ and that ϕ_2 is a fixed smooth weight on X .

Step 1. As a first step, we also assume that $K_1 = X$ and ϕ_1 is smooth. Let μ be a smooth positive volume form, so that both weighted measures (μ, ϕ_i) , $i = 1, 2$ are Bernstein-Markov by Lemma 2.2. By Lemma 2.5, (4.1) is thus equivalent in that case to

$$\lim_{k \rightarrow \infty} \mathcal{L}_k(\mu, \phi_1) - \mathcal{L}_k(\mu, \phi_2) = \mathcal{E}_{\text{eq}}(X, \phi_1) - \mathcal{E}_{\text{eq}}(X, \phi_2). \quad (4.2)$$

As mentioned in (0.9), the volume ratio of L^2 -balls can be expressed as a Gram determinants. As a consequence, we will prove:

Lemma 4.1. *The directional derivatives of $\mathcal{L}_k(\mu, \cdot)$ at a smooth weight ϕ are given by integration against the Bergman measure $\beta(\mu, k\phi)$.*

Proof. Let v be a given smooth function. By (0.9) we have

$$\mathcal{L}_k(\mu, \phi + tv) - \mathcal{L}_k(\mu, \phi) = -\frac{1}{2kN_k} \log \det H_k(t)$$

with the Gram matrix

$$H_k(t) := \left(\int_X s_i^{(k)} \overline{s_j^{(k)}} e^{-2k(\phi + tv)} d\mu \right)_{1 \leq i, j \leq N},$$

$S_k = (s_j^{(k)})_j$ being a fixed orthonormal basis of $H^0(kL)$ with respect to $L^2(\mu, k\phi)$. Since $H_k(0) = \text{id}$, it follows that

$$\begin{aligned} \frac{d}{dt} \mathcal{L}_k(\mu, \phi + tv) &= -\frac{1}{2kN_k} \frac{d}{dt} \text{tr } H_k(t) \\ &= -\frac{1}{2kN_k} \int_X \sum_j |s_j^{(k)}|^2 (-2kv) e^{-2k\phi} \mu = \frac{1}{N_k} \int_X v \rho(\mu, \phi) \mu \end{aligned}$$

and the result follows by definition (2.2) of $\beta(\mu, k\phi)$. $\square$

By Theorem 2.1 we have

$$\lim_{k \rightarrow \infty} \beta(\mu, k\phi) = \mu_{\text{eq}}(X, \phi) \quad (4.3)$$

for any smooth weight ϕ . Now the right-hand side is the derivative of $\mathcal{E}_{\text{eq}}(X, \cdot)$ by Theorem B, so in view of Lemma 4.1 we get (4.2) by integrating (4.3) along the segment between ϕ_1 and ϕ_2 . More precisely, Lemma 4.1 implies

$$\mathcal{L}_k(\mu, \phi_1) - \mathcal{L}_k(\mu, \phi_2) = \int_{t=0}^1 dt \int_X (\phi_1 - \phi_2) \beta(\mu, k\phi_t)$$

with

$$\phi_t := t\phi_1 + (1-t)\phi_2.$$

By dominated convergence (4.3) it follows that

$$\lim_{k \rightarrow \infty} \mathcal{L}_k(\mu, \phi_1) - \mathcal{L}_k(\mu, \phi_2) = \int_{t=0}^1 dt \int_X (\phi_1 - \phi_2) \mu_{\text{eq}}(X, \phi_t)$$

$$= \int_{t=0}^1 \frac{d}{dt} \mathcal{E}_{\text{eq}}(X, \phi_t) = \mathcal{E}_{\text{eq}}(X, \phi_1) - \mathcal{E}_{\text{eq}}(X, \phi_2)$$

by Theorem B, as desired.

Remark 4.2. The argument just presented is similar to Donaldson's proof of Proposition 2 in [Don05a]. In particular, Lemma 4.1 is a variant of Lemma 2 of [Don05a] (cf. also Lemma 3.1 of [Bern06]).

Step 2. We now consider the general case. We first note that

$$\mathcal{L}_k(K, \phi) = \mathcal{L}_k(X, \phi_K)$$

as a consequence of Proposition 1.8, and that $\mathcal{L}_k(X, \cdot)$ is non-decreasing. By Proposition 1.13 we can find two sequences $\phi_j^\pm$ of smooth weights on L such that

$$P_X \phi_j^- \leq \phi_K \leq P_K \phi \leq P_X \phi_j^+ \quad (4.4)$$

where $P_X \phi_j^-$ (resp. $P_X \phi_j^+$) increases (resp. decreases) to $P_K \phi$ almost everywhere (resp. everywhere) on X when j tends to infinity. By Step 1, we get

$$\begin{aligned} \mathcal{E}(P_X \phi_j^-) - \mathcal{E}_{\text{eq}}(X, \phi_2) &= \lim_{k \rightarrow \infty} \mathcal{L}_k(X, P_X \phi_j^-) - \mathcal{L}_k(X, \phi_2) \\ &\leq \liminf_{k \rightarrow \infty} \mathcal{L}_k(X, \phi_K) - \mathcal{L}_k(X, \phi_2) \leq \limsup_{k \rightarrow \infty} \mathcal{L}_k(X, \phi_K) - \mathcal{L}_k(X, \phi_2) \\ &\leq \lim_{k \rightarrow \infty} \mathcal{L}_k(X, P_X \phi_j^+) - \mathcal{L}_k(X, \phi_2) \end{aligned}$$

by (4.4) and monotonicity of $\mathcal{L}_k(X, \cdot)$

$$= \mathcal{E}(P_X \phi_j^+) - \mathcal{E}_{\text{eq}}(X, \phi_2)$$

by Step 1 again. Now

$$\mathcal{E}_{\text{eq}}(X, \phi_j^\pm) = \text{vol}(L)^{-1} \mathcal{E}(P \phi_j^\pm)$$

tends to

$$\text{vol}(L)^{-1} \mathcal{E}(P_K \phi) = \mathcal{E}_{\text{eq}}(K, \phi)$$

by continuity of the energy along monotonic sequences (Proposition 3.3), and putting all this together concludes the proof of Theorem A.

4.2. Proof of Corollary A. Let (μ, ϕ) be a weighted measure with the Bernstein-Markov property and denote by E its support. For each k , we let S_k be an $L^2(\mu, k\phi)$ -orthonormal basis of $H^0(kL)$. Given a weighted subset (K, ψ) , we set

$$\mathcal{D}_k(K, \psi) := \frac{1}{kN_k} \log \|\det S_k\|_{L^\infty(K, k\psi)}$$

with $N_k := h^0(kL)$ and our goal is thus to show that

$$\lim_{k \rightarrow \infty} \mathcal{D}_k(K, \psi) = \mathcal{E}_{\text{eq}}(E, \phi) - \mathcal{E}_{\text{eq}}(K, \psi).$$

As above, we first assume that $K = X$ and ψ is a smooth weight. Let ν be a smooth positive volume form on X , so that (ν, ψ) has the Bernstein-Markov property. By (0.10) from the introduction we have

$$\|\det S_k\|_{L^2(\nu, k\psi)}^2 = N_k! \frac{\text{vol } \mathcal{B}^2(\mu, k\phi)}{\text{vol } \mathcal{B}^2(\nu, k\psi)},$$

or in other words

$$\mathcal{L}_k(\mu, \phi) - \mathcal{L}_k(\nu, \psi) = \frac{1}{kN_k} \log \|\det S_k\|_{L^2(\nu, k\psi)} - \frac{\log N_k!}{2kN_k}.$$

Now $N_k = O(k^n)$ implies

$$\log N_k! = O(k^n \log k) = o(kN_k).$$

Since both (μ, ϕ) and (ν, ψ) are Bernstein-Markov, it thus follows from Theorem A and Lemma 2.5 that

$$\lim_{k \rightarrow \infty} \frac{1}{kN_k} \log \|\det S_k\|_{L^2(\nu, k\psi)} = \mathcal{E}_{\text{eq}}(E, \phi) - \mathcal{E}_{\text{eq}}(X, \psi),$$

and we are therefore reduced to showing that

$$\log \|\det S_k\|_{L^\infty(X, k\psi)} = \log \|\det S_k\|_{L^2(\nu, k\psi)} + o(kN_k).$$

We claim that this follows from the Bernstein-Markov property of (ν, ψ) . Indeed let $\varepsilon > 0$. By the Bernstein-Markov property, there exists $C > 0$ such that

$$|s(x)|_{k\psi}^2 \leq C e^{k\varepsilon} \int_X |s|_{k\psi}^2 d\nu \quad (4.5)$$

for every k , every section $s \in H^0(X, kL)$ and every $x \in X$. Now if $x_1, \dots, x_{N_k}$ are points of X , then for each j

$$x \mapsto \det S_k(x_1, \dots, x_{j-1}, x, x_{j+1}, \dots, x_{N_k})$$

is a holomorphic section in $H^0(X, kL)$. A successive application of (4.5) thus yields

$$\|\det S_k\|_{L^\infty(X, k\phi)}^2 \leq C^{N_k} e^{kN_k\varepsilon} \|\det S_k\|_{L^2(\nu, k\psi)}^2,$$

and the claim follows.

In order to get the general case, we remark that $\mathcal{D}_k(X, \cdot)$ is here again non-decreasing, and a successive application of Proposition 1.8 to each variable of the holomorphic section $\det S_k$ shows that

$$\mathcal{D}_k(K, \psi) = \mathcal{D}_k(X, \psi_K).$$

The same monotone approximation argument as in Step 2 above thus yields the general case of Corollary A.

5. APPLICATIONS TO LOGARITHMIC PLURI-POTENTIAL THEORY

In this section, we will reinterpret our general results in the special case where $(X, L) = (\mathbb{P}^n, \mathcal{O}(1))$ and the compact subsets considered lie in the affine piece $\mathbb{C}^n$. As explained in the introduction, this corresponds to *weighted logarithmic pluri-potential theory* in $\mathbb{C}^n$. For readers primarily interested in this situation we stress that all preliminary results on mixed Monge-Ampère operators in Section 1 are then very standard, cf. for instance [DemBook, DemPP].

We choose homogeneous coordinates $[Z_0 : \dots : Z_n]$ on $\mathbb{P}^n$ such that $Z_0 = 0$ cuts out the hyperplane at infinity, so that $z_j := Z_j/Z_0$ define the euclidian coordinates on $\mathbb{C}^n$. The linear form Z_0 can be seen as the section in $H^0(\mathbb{P}^n, \mathcal{O}(1))$ inducing

the constant polynomial 1 on $\mathbb{C}^n$, and this section enables us to identify *weights* on $\mathcal{O}(1)$ over $\mathbb{C}^n$ to *functions* on $\mathbb{C}^n$ by

$$\phi \mapsto v := \phi - \log |Z_0| = -\log |Z_0|_\phi.$$

We then have $dd^c \phi = dd^c v$ on $\mathbb{C}^n$ by the Lelong-Poincaré formula, and ϕ extends to a psh weight (resp. with minimal singularities) on $\mathcal{O}(1)$ over $\mathbb{P}^n$ iff v is a psh function on $\mathbb{C}^n$ such that $v \leq \log |z| + O(1)$ (resp. $v = \log |z| + O(1)$) at infinity in $\mathbb{C}^n$.

If K is a compact subset of $\mathbb{C}^n$, μ is a probability measure on K and $v \in C^0(K)$ is a continuous function on K , then we will talk about the weighted subset (K, v) and the weighted measure (μ, v) . The equilibrium weight of (K, v) is then identified with the usc regularization of Siciak's extremal function attached to (K, v) , and will be denoted by $P_K v$. It is thus a psh function on $\mathbb{C}^n$ such that $P_K v = \log |z| + O(1)$ at infinity.

5.1. Leja's transfinite diameter as an energy. Denote by $T \subset (\mathbb{C}^*)^n \subset \mathbb{P}^n$ the unit compact torus induced by the toric Kähler structure of $\mathbb{P}^n$. As is well-known and easy to see, the equilibrium function of $(T, 0)$ is then

$$\max_{1 \leq j \leq n} \log^+ |z_j|$$

and the equilibrium measure

$$\mu_T := \mu_{\text{eq}}(T, 0)$$

is then the Haar probability measure on T . For each k , let S_k denote the family of all monomials on $\mathbb{C}^n$ of degree at most k , which is an $L^2(\mu_T, 0)$ -orthonormal basis. Comparing definitions, Leja's transfinite diameter $d_\infty(K, v)$ (cf. [STBook]) is then seen to be defined by

$$\log d_\infty(K, v) = \lim_{k \rightarrow \infty} \frac{(n+1)!}{nk^{n+1}} \log \|\det S_k\|_{L^\infty(K, kv)}$$

provided the limit exists. In the unweighted case ($v = 0$), the limit has been proved to exist by Zaharjuta [Zah75]. Corollary A shows that the limit also exists in the weighted case, and unravelling definitions we get

$$\log d_\infty(K, v) = \frac{1}{n} \sum_{j=0}^n \int_{\mathbb{C}^n} (\max_i \log^+ |z_i| - P_K v) (dd^c P_K v)^j \wedge (dd^c \max_i \log^+ |z_i|)^{n-j}. \quad (5.1)$$

(compare [Rum07, DMR06] for the unweighted case).

5.2. A weighted iterated Robin formula. As a corollary of the recursion formula (3.7) we get the following weighted generalisation of Rumely's Robin-type formula [Rum07]:

Corollary 5.1. *Let (K, v) be a weighted compact subset of $\mathbb{C}^n$. Then its transfinite diameter satisfies*

$$\log d_\infty(K, v) = \frac{1}{n} \sum_{j=0}^n \int_{Y_j} (\log |Z_j| - P_K v) (dd^c P_K v)^{n-j}$$

where $[Z_0 : \cdots : Z_n]$ denote homogeneous coordinates in $\mathbb{P}^n$, $Y_0 = \mathbb{P}^n$ and $Y_j = \{Z_0 = \cdots = Z_{j-1} = 0\}$ when $j \geq 1$.

Proof. Let $T_0 := T$, $\phi_0 = \log |Z_0|$ and $\psi := \log |Z_0| + v$. We then have

$$\frac{n}{n+1} \log d_\infty(K, v) = \mathcal{E}_{Y_0}(P_{T_0}\phi_0) - \mathcal{E}(P_K\psi)$$

with

$$P_T\phi = \max_{0 \leq j \leq n} \log |Z_j|.$$

We thus see that $P_T\phi|_{Y_1}$ coincides with the similarly defined weight $P_{T_1}\phi_1$ on Y_1 . On the other hand $|Z_0|_\phi \equiv 1$ on T and the recursion formula thus implies

$$\begin{aligned} \mathcal{E}_{Y_0}(P_{T_0}\phi_0) - \mathcal{E}(P_K\psi) &= \mathcal{E}_{Y_1}(P_{T_1}\phi_1) - \mathcal{E}_{Y_1}(P_K\psi|_{Y_1}) \\ &\quad + \frac{1}{n+1} \int_{Y_0} (\log |Z_0| - P_K\psi)(dd^c P_K\psi)^n, \end{aligned}$$

and the formula follows by induction. $\square$

In case $n = 1$, this formula relates the *weighted Robin constant*

$$\gamma(K, v) := \lim_{|z| \rightarrow \infty} (v_K^*(z) - \log |z|)$$

to the weighted transfinite diameter by

$$-\log d_\infty(K, v) = \gamma(K, v) + \int_K (P_K v) dd^c(P_K v),$$

the weighted version of Robin's formula (cf. [STBook]).

5.3. Pull-back, the resultant and dynamics. We first consider the following general dynamics situation. Let (X, L) be a projective manifold endowed with an *ample* line bundle and let $f : X \rightarrow X$ be an endomorphism such that $f^*L = dL$ in the Picard group of X for some integer d , called the (first) *algebraic degree* of f . These assumptions imply in particular that f is a finite morphism, and its topological degree is $e = d^n$. We also assume that $d \geq 2$, so that f is not an automorphism.

We would like to consider the action of $d^{-1}f^*$ on the space of weights on L . However the equality $f^*L = dL$ only means that f^*L and dL are isomorphic, and a specific choice of isomorphism is required in order to identify weights on f^*L with weights on dL . Such a choice is equivalent to that of a lifting of f to a map $F : L^* \rightarrow L^*$ that is homogeneous of degree d on the fibres.

The choice of a lift F enables to consider the action of $d^{-1}f^*$ on weights of L , and the *dynamical Green weight* may then be defined by

$$g_F := \lim_{m \rightarrow \infty} (d^{-1}F^*)^m \phi$$

where ϕ is any given continuous psh weight on L . The Green weight g_F is a continuous psh weight, and is the unique fixed point of $d^{-1}F^*$ in the space of continuous weights on L (cf. for instance Sibony's survey [CGSY99]). The Green

weight g_F depends on the specific choice of a lift F and not just on f . Indeed we have

$$g_{\lambda F} = g_F + \frac{1}{d-1} \log |\lambda| \quad (5.2)$$

for each $\lambda \in \mathbb{C}^*$.

Now let (E, ϕ) be a reference weighted subset of X , and define the transfinite diameter (with respect to (E, ϕ)) of a weighted subset (K, ψ) by

$$d_\infty(K, \psi) := \exp \left(\frac{n+1}{n} (\mathcal{E}(P_E \phi) - \mathcal{E}(P_K \phi)) \right),$$

so that this coincides with Leja's transfinite diameter for weighted compact subsets of $\mathbb{C}^n$ if $(E, \phi) = (T, 0)$. We then prove the following general pull-back formula:

Theorem 5.2. *There exists a constant $c > 0$ such that for any weighted subset (K, ψ) we have*

$$d_\infty(f^{-1}K, d^{-1}f^*\psi) = c d_\infty(K, \psi)^{1/d}$$

and in fact

$$c = \exp \left(\frac{(n+1)(d-1)}{nd} (\mathcal{E}(P_E \phi) - \mathcal{E}(g_F)) \right).$$

Proof. Let τ be a psh weight with minimal singularities on L . We have

$$\begin{aligned} \mathcal{E}(P_E \phi) - \mathcal{E}(d^{-1}f^*\tau) &= \mathcal{E}(P_E \phi) - \mathcal{E}(d^{-1}f^*P_E \phi) + \mathcal{E}(d^{-1}f^*P_E \phi) - \mathcal{E}(d^{-1}f^*\tau) \\ &= \mathcal{E}(P_E \phi) - \mathcal{E}(d^{-1}f^*P_E \phi) + d^{-1}(\mathcal{E}(P_E \phi) - \mathcal{E}(\tau)) \end{aligned}$$

by Proposition 3.8. On the other hand Proposition 1.9 shows that the equilibrium weight of $(f^{-1}K, d^{-1}f^*\psi)$ is $d^{-1}f^*P_K\psi$, hence applying this to $\tau := P_K\psi$ proves the first assertion with

$$c := \exp \left(\frac{n+1}{n} (\mathcal{E}(P_E \phi) - \mathcal{E}(d^{-1}f^*P_E \phi)) \right).$$

On the other hand, applying the above relation to $\tau := g_F$ yields

$$\mathcal{E}(P_E \phi) - \mathcal{E}(d^{-1}f^*P_E \phi) = \frac{d-1}{d} (\mathcal{E}(P_E \phi) - \mathcal{E}(g_F)),$$

hence the second assertion. $\square$

We now specialise this transformation formula to $\mathbb{P}^n$ and show how to recover DeMarco-Rumely's result [DMR06].

Corollary 5.3. *Let $f : \mathbb{P}^n \rightarrow \mathbb{P}^n$ be an endomorphism of degree $d \geq 2$, and let $F : \mathbb{C}^{n+1} \rightarrow \mathbb{C}^{n+1}$ be a lifting of f to a d -homogeneous polynomial map. Then for every weighted compact subset (K, ψ) we have*

$$d_\infty(f^{-1}K, d^{-1}f^*\psi) = d_\infty(K, \psi)^{\frac{1}{d}} |\text{Res}(F)|^{-1/nd^{n+1}}$$

where $\text{Res}(F)$ denotes the resultant of F .

Proof. Our arguments mostly follow [BasBer07] and [DMR06] with some simplifications. The space of all d -homogeneous polynomial maps $F : \mathbb{C}^{n+1} \rightarrow \mathbb{C}^{n+1}$ is an affine space $\mathbb{C}^{N+1}$ of dimension

$$N + 1 := (n + 1) \binom{n + d}{d}.$$

Each such map F induces a rational map $f : \mathbb{P}^n \dashrightarrow \mathbb{P}^n$. By [GKZ94] (p.105 and p.427) there exists an *irreducible* hypersurface H of $\mathbb{P}^N$ of degree $(n + 1)d^n$ such that f is an endomorphism iff $F \in \pi^{-1}(\mathbb{P}^N - H)$, where $\pi : \mathbb{C}^{N+1} - \{0\} \rightarrow \mathbb{P}^N$ denotes the quotient map. The variety of all degree d endomorphisms f of $\mathbb{P}^n$ is thus identified with the smooth affine variety $\mathbb{P}^N - H$. The irreducible homogeneous polynomial of degree $(n + 1)d^n$ in $N + 1$ variables cutting out H is called the *resultant* and is denoted by Res . It is normalised by the condition $\text{Res}(F_0) = 1$ for

$$F_0(Z_0, \dots, Z_n) = (Z_0^d, \dots, Z_n^d).$$

The transformation formula (5.2) above implies

$$\mathcal{E}(g_{\lambda F}) = \mathcal{E}(g_F) + \frac{1}{d-1} \log |\lambda|,$$

so that $F \mapsto (d-1)\mathcal{E}(g_F)$ descends to a weight τ on $\mathcal{O}(1)$ over $\mathbb{P}^N - H$.

The main point is now Theorem 4.5 of [BasBer07], which says that $dd^c \tau \equiv 0$ on $\mathbb{P}^N - H$. On the other hand Remark 1.3 of [BasBer07] implies that τ is locally bounded from above near each point of H , hence extends to a psh weight on $\mathcal{O}(1)$ over $\mathbb{P}^N$. The closed positive $(1,1)$ -current $dd^c \tau$ on $\mathbb{P}^N$ is supported on the irreducible hypersurface H , thus the Support Theorem for closed positive currents (see [DemBook] Corollary 2.14 p.165) implies

$$dd^c \tau = c[H]$$

for some $c > 0$, and in fact $c = 1/(n+1)d^n$ since H has degree $(n+1)d^n$. This means in turn that there exists a constant $C > 0$ such that

$$\mathcal{E}(g_F) = \frac{1}{(n+1)(d-1)d^n} \log |\text{Res}(F)| + C \quad (5.3)$$

for all F . This corresponds to Proposition 4.9 of [BasBer07], whose proof has been reformulated here. Now the Green weight of the above map F_0 is easily seen to be

$$g_{F_0} = \max_j \log |Z_j|,$$

which is also the equilibrium weight P_{T0} of $(T, 0)$. Since we have $\text{Res}(F_0) = 1$, we infer that $C = \mathcal{E}_{\text{eq}}(T, 0)$, so that (5.3) becomes

$$\exp \left(\frac{(n+1)(d-1)}{nd} (\mathcal{E}(P_{T0}) - \mathcal{E}(g_F)) \right) = |\text{Res}(F)|^{-1/nd^{n+1}},$$

and the result follows. $\square$

6. ANALYTIC TORSION AND EQUIDISTRIBUTION OF SMALL POINTS

6.1. Asymptotics of the analytic torsion. Let X be a compact Kähler manifold equipped with a fixed Kähler form ω and induced measure ω^n . If L is a line bundle over X , recall that the complex line

$$\det H^\bullet(L) := \sum_{q \geq 0} (-1)^q \det H^q(L)$$

(in our additive notation for tensor products of lines) is called the *determinant of cohomology* of L . If ϕ is a smooth weight on L , then $\det H^\bullet(L)$ can be equipped with a natural L^2 Hermitian metric $|\cdot|_{L^2(\phi)}$, induced by the L^2 metric associated with ϕ and the measure ω^n at the level of harmonic representatives. If ψ is another smooth weight on L , the quotient of the corresponding L^2 metrics on $\det H^\bullet(L)$ yields a number

$$\log \frac{|\cdot|_{L^2(\psi)}^2}{|\cdot|_{L^2(\phi)}^2} = \sum_{q \geq 0} (-1)^q \log \frac{\text{vol } \mathcal{B}_q^2(\phi)}{\text{vol } \mathcal{B}_q^2(\psi)},$$

where we denote by $\mathcal{B}_q^2$ the L^2 -ball of $H^q(X, L)$ for any $q \geq 0$.

The *Ray-Singer analytic torsion* is defined by

$$T(\phi) := \sum_{q \geq 0} (-1)^q q \log \det_{>0} \Delta_q'',$$

where Δ_q'' denotes the anti-holomorphic Laplacian $\overline{\partial}\partial^* + \partial^*\overline{\partial}$ acting on smooth L -valued $(0, q)$ -forms on X , and $\det_{>0}$ denotes the "zeta-regularized" product of its non-zero eigenvalues $0 < \lambda_1 \leq \lambda_2 \leq \dots$, i.e. the derivative at $z = 0$ of the meromorphic continuation to $\mathbb{C}$ of the zeta-function $\sum_j \lambda_j^{-z}$.

The *Quillen metric* on the complex line $\det H^\bullet(L)$ is then the twisted metric

$$|\cdot|_{Q(\phi)}^2 := |\cdot|_{L^2(\phi)}^2 e^{-T(\phi)}.$$

Theorem 1.2.3 of [BGS88] (cf. also [SouBook], Corollary 1 p.132) expresses *variations* of Quillen metrics in terms of secondary Bott-Chern forms. It implies in particular in our case that

$$\sum_{q \geq 0} (-1)^q \log \frac{\text{vol } \mathcal{B}_q^2(\phi)}{\text{vol } \mathcal{B}_q^2(\psi)} + T(\phi) - T(\psi) = \int_X \widetilde{\text{ch}}(\phi, \psi) \wedge \text{td}(\omega) \quad (6.1)$$

for any two smooth weights ϕ, ψ on L , where $\text{td}(\omega) = 1 + \text{Ricci}(\omega)/2 + (\text{higher degree terms})$ is the Todd form of the Hermitian bundle (T_X, ω) and $\widetilde{\text{ch}}$ denotes the secondary form of the Chern character. Formula (3.2) shows that

$$\mathcal{E}(\phi) - \mathcal{E}(\psi) = \frac{n!}{2} \int_X \widetilde{\text{ch}}(\phi, \psi). \quad (6.2)$$

If L is furthermore ample, then the higher cohomology of kL vanishes for $k \gg 1$, thus (6.1) and (6.2) imply

$$\log \frac{\text{vol } \mathcal{B}^2(k\phi)}{\text{vol } \mathcal{B}^2(k\psi)} + T(k\phi) - T(k\psi) = \frac{2k^{n+1}}{n!} (\mathcal{E}(\phi) - \mathcal{E}(\psi)) + O(k^n). \quad (6.3)$$

If ϕ is a smooth weight such that $dd^c\phi > 0$ (hence L is ample), the main result of [BV89] is the following two-term asymptotic expansion of the analytic torsion:

$$\begin{aligned} T(k\phi) &= \frac{1}{2} \int_X \log \frac{(kdd^c\phi)^n}{\omega^n} \exp(kdd^c\phi) + o(k^n) \\ &= \frac{k^n \log k}{2(n-1)!} \text{vol}(L) + \frac{k^n}{2n!} \int_X \log \left(\frac{dd^c\phi}{\omega} \right)^n (dd^c\phi)^n + o(k^n), \end{aligned}$$

and in particular $T(k\phi) = o(k^{n+1})$.

On the other hand, if L is still ample but ϕ has arbitrary curvature, Theorem 10 of [BV89] merely says that $T(k\phi) = O(k^{n+1})$. We will now explain how to refine this estimate using our results:

Theorem 6.1. *Let ω be a Kähler metric on X . If L is an ample line bundle and ϕ is a smooth weight on L with arbitrary curvature, then*

$$\lim_{k \rightarrow \infty} \frac{n!}{2k^{n+1}} T(k\phi) = \mathcal{E}(\phi) - \mathcal{E}(P_X\phi).$$

Proof. Since L is ample, we can choose another smooth weight ψ on L with $dd^c\psi > 0$, so that $T(k\psi) = o(k^{n+1})$ by the result of [BV89] recalled above. On the other hand Lemma 2.5 implies

$$\log \frac{\text{vol } \mathcal{B}^2(k\phi)}{\text{vol } \mathcal{B}^2(k\psi)} = \log \frac{\text{vol } \mathcal{B}^\infty(X, k\phi)}{\text{vol } \mathcal{B}^\infty(X, k\psi)} + o(k^{n+1}),$$

and (6.3) thus yields

$$\log \frac{\text{vol } \mathcal{B}^\infty(X, k\phi)}{\text{vol } \mathcal{B}^\infty(X, k\psi)} + T(k\phi) = \frac{2k^{n+1}}{n!} (\mathcal{E}(\phi) - \mathcal{E}(\psi)) + o(k^{n+1}).$$

Theorem A now yields the result. $\square$

Remark 6.2. We see that for a smooth metric on an ample line bundle Theorem A is in fact *equivalent* to the above estimate for the analytic torsion.

As a consequence of their result on the asymptotics of the analytic torsion, Bismut-Vasserot gave in Theorem 10 of [BV89] an asymptotic comparison result for L^2 metrics induced by two different volume forms. We now give a simple proof of (a generalisation of) that result:

Theorem 6.3. *Let L be a big line bundle and ϕ be an arbitrary smooth weight on L . For any two positive measures μ, ν on X , we then have*

$$\lim_{k \rightarrow \infty} \frac{1}{N_k} \log \frac{\text{vol } \mathcal{B}^2(\nu, k\phi)}{\text{vol } \mathcal{B}^2(\mu, k\phi)} = \int_X \log \left(\frac{\mu}{\nu} \right) \mu_{\text{eq}}(X, \phi).$$

Proof. Note that if f is a function on X we have $\mathcal{B}^2(e^{-f}\mu, \phi) = \mathcal{B}^2(\mu, \phi + 2f)$. Now let $f := \log(\mu/\nu)$ and $\mu_t := e^{-tf}\mu$ for $t \in \mathbb{R}$, so that $\mu_0 = \mu$ and $\mu_1 = \mu'$. By the above remark, Lemma 4.1 implies that

$$\frac{d}{dt} \log \text{vol } \mathcal{B}^2(\mu_t, k\phi) = N_k \int_X f \beta(\mu_t, \phi).$$

We thus get

$$\log \frac{\text{vol } \mathcal{B}^2(\nu, k\phi)}{\text{vol } \mathcal{B}^2(\mu, k\phi)} = N_k \int_{t=0}^1 dt \int_X f \beta(\mu_t, k\phi)$$

and the result follows by dominated convergence since for each t we have $\beta(\mu_t, k\phi) \rightarrow \mu_{\text{eq}}(X, \phi)$ by Theorem 2.1. $\square$

6.2. Adelic heights. Following the discussion in the introduction, let X be a smooth (irreducible) projective variety over $\mathbb{Q}$ and L be a big line bundle on $X/\mathbb{Q}$. Suppose given once and for all an arbitrary collection $(\phi)_p$ of continuous weights on $L_{\mathbb{C}_p}$ over $X(\mathbb{C}_p)$ for each p . If ϕ is a continuous weight on $L_{\mathbb{Q}}$, recall that

$$\mathcal{L}_k^{\mathbb{A}}(\phi) = \frac{1}{kN_k} \log \text{vol}_k \mathcal{B}^{\mathbb{A}}(k\phi)$$

where $\mathcal{B}^{\mathbb{A}}$ denotes the adelic unit-ball defined by (0.12)

By the adelic version of Minkowski's theorem (cf. Appendix A of [BG06]), for every $\varepsilon > 0$ there exists a non-zero $s \in H^0(L)_{\mathbb{Q}}$ such that

$$\log \|s\|_{L^\infty(\phi)} \leq -\mathcal{L}_1^{\mathbb{A}}(\phi) + \log 2 + \varepsilon$$

and $\log \|s\|_{L^\infty(\phi_p)} \leq 0$ for all p .

On the other hand, recall that the *height* of a point $x \in X(\overline{\mathbb{Q}})$ is defined by

$$h_\phi^{\mathbb{A}}(x) := -\frac{1}{\deg(x)} \sum_{y \in Gx} \left(\log |s(y)|_\phi + \sum_p \log |s(y)|_{\phi_p} \right) \quad (6.4)$$

where G denotes the absolute Galois group, Gx is the (finite) Galois orbit of x and s is a rational section of L defined over $\mathbb{Q}$ such that x is neither a pole nor a zero of s . The right-hand side of (6.4) is indeed independent of the choice of s by the product formula, and the sum $\sum_p$ only involves finitely many terms. Note that $h_{k\phi}^{\mathbb{A}}(x) = kh_\phi^{\mathbb{A}}(x)$.

If we use sections $s \in H^0(kL)_{\mathbb{Q}}$ provided by Minkowski's theorem to compute heights, we see by (6.4) that

$$h_\phi^{\mathbb{A}}(x) \geq \mathcal{L}_k^{\mathbb{A}}(\phi) - \frac{\log 2}{k}$$

for any $x \in X(\overline{\mathbb{Q}})$ not in the zero divisor of s , where the adelic $\mathcal{L}$ -functionals $\mathcal{L}_k^{\mathbb{A}}$ are defined by 0.13. As a consequence, if $x_j \in X(\overline{\mathbb{Q}})$ is a *generic* sequence, i.e. a sequence converging to the generic point of X in the Zariski topology, then for each k we get

$$\liminf_j h_\phi^{\mathbb{A}}(x_j) \geq \mathcal{L}_k^{\mathbb{A}}(\phi) - \frac{\log 2}{k},$$

and we infer

$$\liminf_{j \rightarrow \infty} h_\phi^{\mathbb{A}}(x_j) \geq \mathcal{E}_{\text{eq}}^{\mathbb{A}}(\phi). \quad (6.5)$$

Note that we have

$$\mathcal{E}_{\text{eq}}^{\mathbb{A}}(\phi) = \frac{\widehat{\text{vol}}(\overline{L})}{(n+1) \text{vol}(L)}$$

in the notations of [CLT06], p.15, and (6.5) is thus equivalent to Lemma 5.1 of the same [CLT06].

The main point in the proof of Theorem D is the following result.

Lemma 6.4. *The function $\mathcal{E}_{\text{eq}}^{\mathbb{A}}(\cdot)$ is differentiable at any continuous weight ϕ such that $\mathcal{E}_{\text{eq}}^{\mathbb{A}}(\phi) \in \mathbb{R}$. Its directional derivatives are given by integration against the equilibrium measure $\mu_{\text{eq}}(X(\mathbb{C}), \phi)$.*

Proof. Since the Haar measure on

$$H^0(kL)_{\mathbb{A}} \subset H^0(kL)_{\mathbb{R}} \times \Pi_p H^0(kL)_{\mathbb{Q}_p}$$

is induced by a product measure, we see that variations of adelic $\mathcal{L}$ -functionals are given by

$$\mathcal{L}_k^{\mathbb{A}}(\psi) - \mathcal{L}_k^{\mathbb{A}}(\phi) = \frac{1}{kN_k} \log \frac{\text{vol}_k^{\mathbb{R}} \mathcal{B}_{\mathbb{R}}^{\infty}(k\psi)}{\text{vol}_k^{\mathbb{R}} \mathcal{B}_{\mathbb{R}}^{\infty}(k\phi)}$$

where

$$\mathcal{B}_{\mathbb{R}}^{\infty}(\cdot) := \mathcal{B}^{\infty}(\cdot) \cap H^0(kL)_{\mathbb{R}}$$

denotes the unit-ball of the sup-norm on the $\mathbb{R}$ -vector space $H^0(kL)_{\mathbb{R}}$ of $\mathbb{R}$ -sections, and $\text{vol}_k^{\mathbb{R}}$ denotes Lebesgue measure on the latter space. By Lemma 6.5 below, we get

$$\lim_{k \rightarrow \infty} \frac{1}{kN_k} \log \frac{\text{vol}_k^{\mathbb{R}} \mathcal{B}_{\mathbb{R}}^{\infty}(k\psi)}{\text{vol}_k^{\mathbb{R}} \mathcal{B}_{\mathbb{R}}^{\infty}(k\phi)} = \lim_{k \rightarrow \infty} \frac{1}{2kN_k} \log \frac{\text{vol}_k \mathcal{B}^{\infty}(k\psi)}{\text{vol}_k \mathcal{B}^{\infty}(k\phi)},$$

where $\mathcal{B}^{\infty}$ denotes as before the unit-ball of the sup-norm in the *complex* vector space $H^0(kL)_{\mathbb{C}}$ of $\mathbb{C}$ -sections and vol_k is Lebesgue measure on that space. We now conclude by Theorems A and B, using the trivial relation

$$\limsup_{k \rightarrow \infty} a_k - \limsup_{k \rightarrow \infty} b_k = \lim_{k \rightarrow \infty} (a_k - b_k)$$

provided the right-hand limit exists (and is finite). $\square$

Lemma 6.5. *Let X be a smooth projective variety defined over $\mathbb{R}$, and let L be a big line bundle on $X/\mathbb{R}$. Let ϕ be a continuous weight over $X(\mathbb{C})$, and denote by $\mathcal{B}_{\mathbb{R}}^{\infty}(\phi)$ the unit-ball of the sup-norm in $H^0(L)_{\mathbb{R}}$. Then*

$$\log \frac{\text{vol}_k^{\mathbb{R}} (\mathcal{B}_{\mathbb{R}}^{\infty}(k\phi))^2}{\text{vol} \mathcal{B}^{\infty}(k\phi)} = o(kN_k).$$

Proof. Let μ be a smooth positive volume form on $X(\mathbb{C})$, so that (μ, ϕ) has the Bernstein-Markov property. The easy scaling argument used in the proof of Lemma 2.5 immediately yields

$$\log \frac{\text{vol}_k^{\mathbb{R}} (\mathcal{B}_{\mathbb{R}}^2(k\phi))^2}{\text{vol}_k \mathcal{B}^2(k\phi)} = \log \frac{\text{vol}_k^{\mathbb{R}} (\mathcal{B}_{\mathbb{R}}^{\infty}(k\phi))^2}{\text{vol}_k \mathcal{B}^{\infty}(k\phi)} + o(kN_k).$$

But we can further assume that μ is invariant by complex conjugation, so that the $L^2(\mu, k\phi)$ -scalar product is defined over $\mathbb{R}$, and it is then easy to see that the left-hand side is equal to its value in the Euclidian space situation, that is

$$\frac{\text{vol}_k^{\mathbb{R}} (\mathcal{B}_{\mathbb{R}}^2(k\phi))^2}{\text{vol} \mathcal{B}^2(k\phi)} = \frac{N_k!}{((N_k/2)!)^2}$$

by expressing it in terms of Gram determinants of orthonormal basis of $H^0(L)_{\mathbb{R}}$. Now both N_k and $N_k/2$ are $O(k^n)$, and this implies by Stirling's formula that both $\log N_k!$ and $\log(N_k/2)!$ are $O(k^n \log k) = o(k^{n+1})$. The result follows. $\square$

6.3. Proof of Theorem C. If $x \in X(\overline{\mathbb{Q}})$ is an algebraic point, let μ_x denote the averaging measure on $X(\mathbb{C})$ along the Galois orbit Gx . By (6.4) it is immediate to see that

$$h_{\phi+v}^{\mathbb{A}}(x) = h_{\phi}^{\mathbb{A}}(x) + \langle \mu_x, v \rangle \quad (6.6)$$

for any continuous function v on $X(\mathbb{C})$.

Now let (x_j) be a generic sequence such that $\lim_{j \rightarrow \infty} h_{\phi}(x_j) = \mathcal{E}_{\text{eq}}^{\mathbb{A}}(\phi) \in \mathbb{R}$. If v is a continuous function on $X(\mathbb{C})$, we are to show that

$$\lim_{j \rightarrow \infty} \langle \mu_{x_j}, v \rangle = \langle \mu_{\text{eq}}(X(\mathbb{C}), \phi), v \rangle.$$

By Lemma 6.4 the right-hand side is equal to the derivative at $t = 0$ of the function $g(t) := \mathcal{E}_{\text{eq}}^{\mathbb{A}}(\phi + tu)$. On the other hand by (6.6) the left-hand side is equal to the derivative at $t = 0$ of the affine function $f_j(t) := h_{\phi+tu}(x_j)$. The asymptotic lower bound (6.5) implies that

$$\liminf_{j \rightarrow \infty} f_j(t) \geq g(t)$$

for all t , and the following elementary lemma yields the result.

Lemma 6.6. *Let f_j be a sequence of concave functions on $\mathbb{R}$ and let g be a function on $\mathbb{R}$ such that*

- $\liminf_{j \rightarrow \infty} f_j \geq g$.
- $\lim_{j \rightarrow \infty} f_j(0) = g(0)$.

If the f_j and g are differentiable at 0, then

$$\lim_{j \rightarrow \infty} f_j'(0) = g'(0).$$

Proof. Since f_j is concave, we have

$$f_j(0) + f_j'(0)t \geq f_j(t)$$

and it follows that

$$\liminf_{j \rightarrow \infty} t f_j'(0) \geq g(t) - g(0).$$

The result now follows by first letting $t > 0$ and then $t < 0$ tend to 0. $\square$

This lemma is inspired by the variational principle in the original proof by Szpiro-Ullmo-Zhang [SUZ97]. The case of concave functions f_k pertains to the situation considered in [BBWN08].

REFERENCES

- [AB95] Abbes, A.; Bouche, T.: Théorème de Hilbert-Samuel “arithmétique”. Ann. Inst. Fourier 45 (1995), no. 2, 375–401.
- [Ale38] Alexandrov A.D.: On the theory of mixed volumes of convex bodies III: Extension of two theorems of Minkowski on convex polyhedra to arbitrary convex bodies (Russian). Mat. Sbornik 3 (1938), no.1, 27–44. [English translation available in Selected works part I: Selected scientific papers. Gordon and Breach]

- [Aub84] Aubin, T.: Réduction du cas positif de l'équation de Monge-Ampère sur les variétés kählériennes compactes à la démonstration d'une inégalité. *J. Funct. Anal.* 57 (1984), no. 2, 143–153.
- [BasBer07] Bassanelli, G.; Bertolotto, F.: Bifurcation currents in holomorphic dynamics on $\mathbb{P}^n$. *J. Reine. Angew. Math* 608 (2007), 201–235
- [Ber06] Berman, R.: Super Toeplitz operators on holomorphic line bundles *J. Geom. Anal.* 16 (2006), no. 1, 1–22.
- [Ber07a] Berman, R.: Bergman kernels and weighted equilibrium measures of $\mathbb{C}^n$. Preprint (2007) arXiv:math/0702357. To appear in *Indiana Univ. Journ. of Math.*
- [Ber07b] Berman, R.: Bergman kernels and equilibrium measures for line bundles over projective manifolds. Preprint (2007) arXiv:0710.4375.
- [BerBou08a] Berman, R.; Boucksom, S.: Capacities and weighted volumes for line bundles. Preprint (2008) arXiv:0803.1950.
- [BerBou08b] Berman, R.; Boucksom, S.: Equidistribution of Fekete points on complex manifolds. Preprint (2008) arXiv:0807.0035.
- [BBWN08] Berman, R.; Boucksom, S.; Witt Nyström, D.: Convergence towards equilibrium on complex manifolds. Preprint.
- [BWN08] Berman, R.; Witt Nyström, D.: Convergence of Bergman measures for high powers of a line bundle. Preprint (2008) arXiv:0805.2846.
- [Bern06] Berndtsson, B.: Positivity of direct image bundles and convexity on the space of Kähler metrics. Preprint (2006) arXiv.math.CV/0608385. To appear in *J. Diff. Geom.*
- [BGS88] Bismut, J-M; Gillet, H.; Soulé, C.: Analytic torsion and holomorphic determinant bundles. I-III. Direct images and Bott-Chern forms. *Comm. Math. Phys.* 115 (1988).
- [BV89] Bismut, J.-M.; Vasserot, E.: The asymptotics of the Ray-Singer analytic torsion associated with high powers of a positive line bundle. *Comm. Math. Phys.* 125 (1989), no. 2, 355–367
- [Blo06] Bloom, T.: Weighted polynomials and weighted pluripotential theory. arXiv:math/0610330. To appear in *Trans. A.M.S.*
- [BBLW08] Bloom, T.; Bos, L.; Levenberg, N.; Waldron, S.: On the convergence of optimal measures. Preprint (2008) arXiv:0808.0762.
- [BC99] Bloom, T.; Calvi, J.-P., On the multivariate transfinite diameter. *Ann. Polon. Math.* 72 (1999), no. 3, 285–305.
- [BL03] Bloom, T.; Levenberg, N.: Weighted pluripotential theory in $\mathbb{C}^N$ *Amer. J. Math.* 125 (3) (2003), 57–103.
- [BL07a] Bloom, T.; Levenberg, N.: Strong asymptotics for Christoffel functions of planar measures. Preprint (2007) arXiv: 0709.2073.
- [BL07b] Bloom, T.; Levenberg, N.: Transfinite diameter notions in $\mathbb{C}^N$ and integrals of Vandermonde determinants. Preprint (2007) arXiv: 0712.2844.
- [BG06] Bombieri, E.; Gubler, W.: Heights in Diophantine geometry, *New Mathematical Monographs*, 4. Cambridge University Press, Cambridge, 2006.
- [Bon98] Bonavero, L.: Inégalités de Morse holomorphes singulières. *J. Geom. Anal.* 8 (1998), no. 3, 409–425.
- [BGS94] Bost, J.-B.; Gillet, H.; Soulé, C.: Heights of projective varieties and positive Green forms. *J. Amer. Math. Soc.* 7 (1994), no. 4, 903–1027.
- [Bo90] Bouche, T.: Convergence de la métrique de Fubini-Study d'un fibré linéaire positif, *Ann. Inst. Fourier* (1) 40 (1990), 117–130.
- [Bou02] Boucksom, S.: On the volume of a line bundle. *Internat. J. Math* 13 (2002), no 10. 1043–1063
- [Bou04] Boucksom, S.: Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. Ecole Norm. Sup.* 37 (2004), 45–76

- [BDPP04] Boucksom, S.; Demailly, J.-P.; Paun, M.; Peternell, T.: The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. arXiv:math/0405285.
- [BEGZ08] Boucksom, S.; Eyssidieux, P.; Guedj, V.; Zeriahi, A.: Monge-Ampère equations in big cohomology classes. Preprint (2008) arXiv:0812.3674.
- [BFJ07] Boucksom, S.; Favre, C.; Jonsson, M.: Differentiability of volumes of divisors and a problem of Teissier. To appear in *J. of Algebraic Geom.*
- [BFJ08] Boucksom, S.; Favre, C.; Jonsson, M.: Valuations and plurisubharmonic singularities. *Publ. Res. Inst. Math. Sci.* 44 (2008), no. 2, 449–494.
- [Cat99] Catlin, D.: The Bergman kernel and a theorem of Tian. In *Analysis and geometry in several complex variables* (Katata, 1997), Trends Math., pages 1-23. Birkhäuser Boston, Boston, MA, 1999.
- [CGSY99] Cerveau, D.; Ghys, E.; Sibony, N.; Yoccoz, J.-C.: Dynamique et géométrie complexes. *Panoramas et Synthèses* 8 (1999). Société Mathématique de France, Paris.
- [CLT06] Chambert-Loir, A.; Thuillier, A.: Formule de Mahler et équidistribution logarithmique. Preprint (2006) arXiv:math.NT/0612556.
- [Che00] Chen, X.X.: The space of Kähler metrics, *J. Differential Geom.* 56 (2000), no. 2, 189-234.
- [Dei99] Deift, P.A.: Orthogonal polynomials and random matrices: a Riemann-Hilbert approach. *Courant Lecture Notes in Mathematics*, 3. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999.
- [DemBook] Demailly, J.-P.: Complex analytic and algebraic geometry. Book available at www-fourier.ujf-grenoble.fr/~demailly/books.html.
- [DemPP] Demailly, J.-P.: Potential Theory in Several Complex Variables. Manuscript available at www-fourier.ujf-grenoble.fr/~demailly/.
- [Dem92] Demailly, J.-P.: Regularization of closed positive currents and Intersection Theory. *J. Alg. Geom.* 1, (1992), 361-409.
- [DMR06] De Marco, L.; Rumely, R.: Transfinite diameter and the resultant. arXiv:math/0601109.
- [Don85] Donaldson, S.K.: Anti self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles. *Proc. London Math. Soc.* (3) 50 (1985), no. 1, 1–26.
- [Don05a] Donaldson, S.K.: Scalar curvature and projective embeddings. II. *Q. J. Math.* 56 (2005), no. 3, 345–356.
- [Don05b] Donaldson, S.K.: Some numerical results in complex differential geometry. Preprint (2005) arXiv:math/0512625.
- [GZ05] Guedj, V.; Zeriahi, A.: Intrinsic capacities on compact Kähler manifolds. *J. Geom. Anal.* 15 (2005), 607-639.
- [GKZ94] Gelfand, I. M.; Kapranov, M. M.; Zelevinsky, A.V.: Discriminants, resultants, and multidimensional determinants. Birkhäuser Boston, Inc., Boston, MA, 1994. x+523.
- [JS93] Ji, S.; Shiffman, B.: Properties of compact complex manifolds carrying closed positive currents. *J. Geom. Anal.* 3 (1993), no. 1, 37–61.
- [KliBook] Klimek, M.: Pluripotential theory. *London Mathematical Society Monographs. New Series*, 6. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1991.
- [LazBook] Lazarsfeld, R.: Positivity in algebraic geometry. I. Classical setting: line bundles and linear series. II. Positivity for vector bundles, and multiplier ideals. *A Series of Modern Surveys in Mathematics*, 48. and 49. Springer-Verlag, Berlin, 2004.
- [LM80] Lempio, F.; Maurer, H.: Differential stability in infinite-dimensional nonlinear programming. *Appl. Math. Optim.* 6 (1980), 139–152.
- [Mab86] Mabuchi, T.: K-energy maps integrating Futaki invariants. *Tohoku Math. J.* (2) 38 (1986), no. 4, 575–593.

- [PS03] Phong, D.H.; Sturm, J.: Stability, energy functionals, and Kähler-Einstein metrics. *Comm. Anal. Geom.* 11 (2003), no. 3
- [RokBook] Rockafellar, R.T.: *Convex analysis*. Princeton Landmarks in Mathematics, 1970.
- [Rum07] Rumely, R.: A Robin formula for the Fekete-Leja transfinite diameter. *Math. Ann.* 337 (2007), no. 4, 729–738.
- [RLV00] Rumely, R.; Lau, C.F.; Varley, R.: Existence of the sectional capacity. *Mem. Amer. Math. Soc.* 145 (2000), no. 690.
- [STBook] Saff, E.B.; Totik, V.: *Logarithmic potentials with exterior fields*. Springer-Verlag, Berlin. (1997) (with an appendix by Bloom, T.).
- [SchBook] Schneider, R.: *Convex bodies, the Brunn-Minkowski theory*. Volume 44 of *Encyclopaedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1993.
- [Sic88] Siciak, J.: Families of polynomials and determining measures. *Ann. Fac. Sci. Toulouse Math.* (5) 9 (1988), no. 2, 193–211.
- [SouBook] Soulé, C.: *Lectures on Arakelov geometry*. With the collaboration of D. Abramovich, J.-F. Burnol and J. Kramer. *Cambridge Studies in Advanced Mathematics* 33, 1992.
- [SUZ97] Szpiro, L.; Ullmo, E.; Zhang, S.: Equirépartition des petits points. *Invent. Math.* 127 (1997), no. 2, 337–347.
- [Tia90] Tian, G.: On a set of polarized Kähler metrics on algebraic manifolds. *J. Differential Geom.* 32 (1) (1990), 99–130.
- [TiaBook] Tian, G.: *Canonical metrics in Kähler geometry*. Birkhäuser (2000).
- [Tia00] Tian, G.: Bott-Chern forms and geometric stability. *Discrete Contin. Dynam. Systems* 6 (2000), no.1, 211–220.
- [Yua06] Yuan, X.: Big line bundles over arithmetic varieties. Preprint (2006) arXiv:math/0612424. To appear in *Invent. Math.*
- [Zah75] Zaharjuta, V.: Transfinite diameter, Chebyshev constants, and capacity for compacta in $\mathbb{C}^n$, *Math. USSR Sbornik* 25 (1975), 350–364.
- [Zel98] Zelditch, S.: Szegő kernels and a theorem of Tian. *Int. Math. Res. Notices* 6 (1998), 317–331.
- [Zha06] Zhang, S.: Distributions in algebraic dynamics. A tribute to Professor S.S. Chern, *Surveys in Differential Geometry*, vol 10, 381–430, International Press 2006 (2006).

UNIVERSITÉ GRENOBLE I, INSTITUT FOURIER, SAINT-MARTIN D'HÈRES, FRANCE

E-mail address: robertb@math.chalmers.se

CNRS-UNIVERSITÉ PARIS 7, INSTITUT DE MATHÉMATIQUES, F-75251 PARIS CEDEX 05, FRANCE

E-mail address: boucksom@math.jussieu.fr